

# The quasiclassical limit of the symmetry constraint of the KP hierarchy and the dispersionless KP hierarchy with self-consistent sources

Ting XIAO and Yunbo ZENG

*Department of Mathematical Sciences, Tsinghua University, Beijing 100084, P.R.China*  
*E-mail: xiaoting97@mails.tsinghua.edu.cn, yzeng@math.tsinghua.edu.cn*

*Received May 13, 2005; Accepted September 28, 2005*

## Abstract

For the first time we show that the quasiclassical limit of the symmetry constraint of the Sato operator for the KP hierarchy leads to the generalized Zakharov reduction of the Sato function for the dispersionless KP (dKP) hierarchy which has been proved to be result of symmetry constraint of the dKP hierarchy recently. By either regarding the symmetry constrained dKP hierarchy as its stationary case or taking the dispersionless limit of the KP hierarchy with self-consistent sources directly, we construct a new integrable dispersionless hierarchy, i.e., the dKP hierarchy with self-consistent sources and find its associated conservation equations (or equations of Hamilton-Jacobi type). Some solutions of the dKP equation with self-consistent sources are also obtained by hodograph transformations.

## 1 Introduction

The dispersionless integrable hierarchies provide us an interesting type of nonlinear integrable models which have important applications from complex analysis to topological field theory (see [1-16]). In [14, 25, 6, 7, 22], a standard procedure of dispersionless limit of integrable dispersionfull hierarchies is proposed. In this procedure, dispersionless hierarchies arise as the quasiclassical limit of the original dispersionfull Lax equations performed by replacing operators by phase space functions, commutators by Poisson brackets and the role of Lax pair equations by conservation equations (or equations of Hamilton-Jacobi type). A  $\bar{\partial}$  scheme of dispersionless hierarchies has been proposed by Konopelchenko et al in [13-16]. Recently, from this point of view, some important reductions of dispersionless hierarchies are shown to be nothing but results of symmetry constraints [3]. Also several methods for solving dispersionless hierarchies have been formulated such as twistorial method [21, 22, 16] and hodograph transformation [6, 7]. In [6, 7], from the conservation equations of the dispersionless KP equation, Kodama and Gibbons found exact solutions of it and its reductions by hodograph transformations and obtained general hodograph

---

equations for hydrodynamical type equations.

The soliton equations with self-consistent sources (SESCS) are another type of important integrable models in many fields of physics, such as hydrodynamics, state physics, plasma physics, etc (see [18-24]). For example, the KP equation with self-consistent sources (KPESCS) describes the interaction of a long wave with a short-wave packet propagating on the  $x, y$  plane at an angle to each other (see [17] and the references therein). We proposed a general approach to construct SESCO by regarding the constrained integrable hierarchy as stationary system of the corresponding hierarchy with self-consistent sources, and found the Lax representation for the latter naturally from that for the former [21-24]. In this sense, the soliton hierarchies with self-consistent sources may be viewed as integrable generalizations of the original soliton hierarchies.

In contrast with the dispersionfull integrable hierarchies with self-consistent sources, the dispersionless integrable hierarchies with self-consistent sources have not been studied yet. In this paper, we first investigate the quasiclassical limit of the symmetry constraint of the Sato operator for the KP hierarchy which leads to the generalized Zakharov reduction of the Sato function for the dKP hierarchy. The latter has recently been proved to be result of symmetry constraint of the dKP hierarchy using the  $\bar{\partial}$  method by Bogdanov and Konopelchenko [3]. So, we find the relation that the symmetry constraint of the Sato function for the dispersionless hierarchy can be obtained by the quasiclassical limit of the symmetry constraint of the Sato operator for the corresponding dispersionfull hierarchy. The key point for the quasiclassical limit here is to use the asymptotic forms of the Baker and adjoint Baker functions and regard the quantities  $q_i$  and  $r_i$  as Baker and adjoint Baker functions when the spectral parameter  $\lambda$  evaluated at some points  $\lambda_i$ . By either regarding the symmetry constrained dKP hierarchy as its stationary case or taking the dispersionless limit of the KP hierarchy with self-consistent sources directly, we construct a new dispersionless hierarchy, i.e., the dKP hierarchy with self-consistent sources. This hierarchy is also integrable since we can find its associated conservation equations (or equations of Hamilton-Jacobi type). So in this sense, the dKP hierarchy with self-consistent sources may be viewed as an integrable generalization of the dKP hierarchy. Compared with the case of a nonlocal term appearing in the  $t$ -part of the Lax pair for the KP equation with self-consistent sources [23], the  $t$ -part of the conservation equations for the dKP equation with self-consistent sources possesses rational terms with poles. From the obtained conservation equations, we can solve the dKP hierarchy with self-consistent sources by hodograph transformations.

The paper will be organized as follows. In section 2, we briefly review some definitions and results about the KP hierarchy with self-consistent sources. In section 3, we take the quasiclassical limit to the symmetry constraint of the Sato operator for the KP hierarchy to obtain the symmetry constraint of the Sato function for the dKP hierarchy. The dKP hierarchy with self-consistent sources and its associated conservation equations are obtained. Section 4 is devoted to solving the dKP hierarchy with self-consistent sources by hodograph transformations and some solutions for the dKP equation with self-consistent sources are presented.

## 2 The KP hierarchy with self-consistent sources

We first review some definitions and results about the KP hierarchy with self-consistent sources in the framework of Sato theory [23-26]. Given a pseudo-differential operator (PDO) of the form

$$L = \partial + u_1(t)\partial^{-1} + u_2(t)\partial^{-2} + \dots, \quad (2.1)$$

where  $\partial = \partial_x$ ,  $\partial\partial^{-1} = \partial^{-1}\partial = 1$ ,  $t = (t_1 = x, t_2, \dots)$ , the KP hierarchy is defined as

$$\partial_{t_n} L = [B_n, L], \quad (2.2)$$

where  $B_n = (L^n)_{\geq 0}$  is the differential part of  $L^n$ . The Lax equation (2.2) is equivalent to the existence of the Baker function  $\psi$  such that

$$L\psi = \lambda\psi, \quad (2.3a)$$

$$\partial_{t_n}\psi = B_n\psi, \quad (2.3b)$$

and  $\psi$  also satisfies

$$\partial_\lambda\psi = M\psi, \quad (2.4)$$

where  $M$  is the Orlov operator of the KP hierarchy. The adjoint Baker function  $\psi^*$  satisfies

$$L^*\psi^* = \lambda\psi^*, \quad (2.5a)$$

$$\partial_n\psi^* = -B_n^*\psi^*, \quad (2.5b)$$

$$\partial_\lambda\psi^* = -M^*\psi^*. \quad (2.5c)$$

Making a constraint of the PDO  $L$  (2.1) as [19]

$$L^n = (L^n)_{\geq 0} + \sum_1^N q_i(t)\partial^{-1}r_i(t), \quad n \in \mathbb{N} \quad (2.6)$$

where  $q_i(t)$  and  $r_i(t)$  satisfying

$$q_{i,t_m} = B_m q_i, \quad r_{i,t_m} = -B_m^* r_i, \quad B_m = [(L^n)^{\frac{m}{n}}]_{\geq 0}, \quad i = 1, \dots, N, \forall m \in \mathbb{N}, \quad (2.7)$$

we will get the  $n$ -constrained KP hierarchy as

$$(L^n)_{t_m} = [B_m, L^n], \quad (2.8a)$$

$$q_{i,t_m} = B_m q_i, \quad (2.8b)$$

$$r_{i,t_m} = -B_m^* r_i, \quad i = 1, \dots, N. \quad (2.8c)$$

If we add the term  $(B_m)_{t_n}$  to the right side of equation (2.8a), the KP hierarchy with self-consistent sources will be obtained as [23]

$$(B_m)_{t_n} - (L^n)_{t_m} + [B_m, L^n] = 0, \quad (2.9a)$$

$$q_{i,t_m} = B_m q_i, \tag{2.9b}$$

$$r_{i,t_m} = -B_m^* r_i, \quad i = 1, \dots, N. \tag{2.9c}$$

As many cases in (1+1)-dimension (see [27, 28] and the references therein), the  $n$ -constrained KP hierarchy may be considered as the stationary one of the KP hierarchy with self-consistent sources if " $t_n$ " is viewed as the evolution variable and the Lax representation for the KP hierarchy with self-consistent sources can be obtained naturally from that for the constrained KP hierarchy [23]. The most important equation in the hierarchy is the KP equation with self-consistent sources obtained when  $n = 3, m = 2$  in (2.9), [17, 18]

$$[u_{1,t} - 3u_1 u_{1,x} - \frac{1}{4}u_{1,xxx} + \sum_{i=1}^N (q_i r_i)_x]_x - \frac{3}{4}u_{1,yy} = 0, \tag{2.10a}$$

$$q_{i,y} = q_{i,xx} + 2u_1 q_i, \tag{2.10b}$$

$$r_{i,y} = -r_{i,xx} - 2u_1 r_i, \quad i = 1, \dots, N \tag{2.10c}$$

where  $t = t_3, y = t_2$ . With (2.10b) and (2.10c), (2.10a) will be obtained by the compatibility of the following linear equations (Lax pair)[23]

$$\psi_y = \psi_{xx} + 2u_1 \psi, \tag{2.11a}$$

$$\psi_t = \psi_{xxx} + 3u_1 \psi_x + \frac{3}{2}(u_{1,x} + (\partial^{-1}u_{1,y}))\psi + \sum_{i=1}^N q_i \partial^{-1}(r_i \psi). \tag{2.11b}$$

### 3 Dispersionless limit

Following the standard procedure of dispersionless limit introduced in [14, 25, 6, 7, 22], we will get the dispersionless counterpart of (2.9) which can be regarded as the dispersionless KP hierarchy with self-consistent sources. Simply taking  $T_n = \epsilon t_n$  and thinking of  $u_n(\frac{T}{\epsilon}) = U_n(T) + O(\epsilon)$  as  $\epsilon \rightarrow 0, L$  in (2.1) changes into

$$L_\epsilon = \epsilon \partial + \sum_{i=1}^{\infty} u_i(\frac{T}{\epsilon})(\epsilon \partial)^{-i} = \epsilon \partial + \sum_{i=1}^{\infty} (U_i(T) + O(\epsilon))(\epsilon \partial)^{-i}, \quad \partial = \partial_X, \quad X = \epsilon x. \tag{3.1}$$

The constraint (2.6) now changes into

$$L_\epsilon^n = B_{\epsilon n} + \sum_{i=1}^N q_i(\frac{T}{\epsilon})(\epsilon \partial)^{-1} r_i(\frac{T}{\epsilon}), \quad B_{\epsilon n} = (L_\epsilon^n)_{\geq 0}, \tag{3.2}$$

where  $q_i(\frac{T}{\epsilon})$  and  $r_i(\frac{T}{\epsilon})$  satisfy

$$\epsilon [q_i(\frac{T}{\epsilon})]_{T_m} = B_{\epsilon m} q_i(\frac{T}{\epsilon}), \quad \epsilon [r_i(\frac{T}{\epsilon})]_{T_m} = -B_{\epsilon m}^* r_i(\frac{T}{\epsilon}), \quad B_{\epsilon m} = [(L_\epsilon^n)^{\frac{m}{n}}]_{\geq 0}, \quad i = 1, \dots, N, \tag{3.3}$$

and the counterpart of (2.8) is

$$\epsilon (L_\epsilon^n)_{T_m} = [B_{\epsilon m}, L_\epsilon^n], \tag{3.4a}$$

$$\epsilon[q_i(\frac{T}{\epsilon})]_{T_m} = B_{\epsilon m} q_i(\frac{T}{\epsilon}), \quad (3.4b)$$

$$\epsilon[r_i(\frac{T}{\epsilon})]_{T_m} = -B_{\epsilon m}^* r_i(\frac{T}{\epsilon}), \quad i = 1, \dots, N. \quad (3.4c)$$

It is proved in [22] that

$$\mathcal{L} = \sigma^\epsilon(L_\epsilon) = p + \sum_{i=1}^{\infty} U_i(T) p^{-i}. \quad (3.5)$$

is a solution of the dKP hierarchy, i.e., satisfies

$$\partial_{T_n} \mathcal{L} = \{\mathcal{B}_n, \mathcal{L}\}, \quad (3.6)$$

where  $\sigma^\epsilon$  denotes the principal symbol [22], the Poisson bracket is defined as

$$\{A(p, x), B(p, x)\} = \frac{\partial A}{\partial p} \frac{\partial B}{\partial x} - \frac{\partial A}{\partial x} \frac{\partial B}{\partial p}, \quad (3.7)$$

and  $\mathcal{B}_n = (\mathcal{L})_{\geq 0}^n$  now refers to powers of  $p$ .

The dKP hierarchy can be also written in the zero-curvature form

$$\frac{\partial \mathcal{B}_n}{\partial T_m} - \frac{\partial \mathcal{B}_m}{\partial T_n} + \{\mathcal{B}_n, \mathcal{B}_m\} = 0. \quad (3.8)$$

In [22], from (2.3) and (2.4) (with  $L$ ,  $B_n$ ,  $M$  and  $\partial_n$  replaced by  $L_\epsilon$ ,  $B_{\epsilon n}$ ,  $M_\epsilon$  and  $\epsilon \partial_{T_n}$  respectively), it has been proved that  $\psi(\frac{T}{\epsilon})$  has the following WKB asymptotic expansion as  $\epsilon \rightarrow 0$

$$\psi(\frac{T}{\epsilon}) = \exp[\frac{1}{\epsilon} S(T, \lambda) + O(1)], \quad \epsilon \rightarrow 0. \quad (3.9)$$

In order to take the quasiclassical limit of the constraint (3.2), we also need to find the asymptotic form of the adjoint Baker function. Similarly like the proof in [22], from (2.5), we can find  $\psi^*(\frac{T}{\epsilon})$  has the following WKB asymptotic expansion

$$\psi^*(\frac{T}{\epsilon}) = \exp[-\frac{1}{\epsilon} S(T, \lambda) + O(1)], \quad \epsilon \rightarrow 0. \quad (3.10)$$

From (2.3b) and (3.9), we obtain a hierarchy of conservation equations for the momentum function  $p = \frac{\partial S}{\partial X}$ ,

$$\frac{\partial p}{\partial T_n} = \frac{\partial \mathcal{B}_n(p)}{\partial X}, \quad (3.11)$$

the compatibility of which, i.e.,  $\frac{\partial^2 p}{\partial T_n \partial T_m} = \frac{\partial^2 p}{\partial T_m \partial T_n}$  implies the dKP hierarchy (3.8).

Regarding

$$q_i(\frac{T}{\epsilon}) = \psi(\frac{T}{\epsilon}, \lambda = \lambda_i) \sim \exp[\frac{S(T, \lambda_i)}{\epsilon} + \alpha_{i1} + O(\epsilon)], \quad \epsilon \rightarrow 0, \quad (3.12a)$$

$$r_i(\frac{T}{\epsilon}) = \psi^*(\frac{T}{\epsilon}, \lambda = \lambda_i) \sim \exp[-\frac{S(T, \lambda_i)}{\epsilon} + \alpha_{i2} + O(\epsilon)], \quad \epsilon \rightarrow 0, \quad i = 1, \dots, N, \quad (3.12b)$$

we will find that

$$q_i\left(\frac{T}{\epsilon}\right)(\epsilon\partial)^{-1}r_i\left(\frac{T}{\epsilon}\right) = e^{\alpha_{i1}+\alpha_{i2}}[(\epsilon\partial)^{-1} + \left(\frac{\partial S(T,\lambda_i)}{\partial X}\right) + O(\epsilon)](\epsilon\partial)^{-2} + \left[\left(\frac{\partial S(T,\lambda_i)}{\partial X}\right)^2 + O(\epsilon)\right](\epsilon\partial)^{-3} + \dots], \quad \epsilon \rightarrow 0.$$

Taking the principal symbol of both sides of (3.2), we have

$$\begin{aligned} \mathcal{L}^n &= \mathcal{B}_n + \sum_{i=1}^N e^{\alpha_{i1}+\alpha_{i2}} [p^{-1} + S_X(T, \lambda_i)p^{-2} + S_X^2(T, \lambda_i)p^{-3} + \dots] \\ &= \mathcal{B}_n + \sum_{i=1}^N \frac{v_i}{p - p_i} \end{aligned} \quad (3.13)$$

where  $\mathcal{B}_n = (\mathcal{L}^n)_{\geq 0}$  and

$$v_i = e^{\alpha_{i1}+\alpha_{i2}}, \quad p_i = S_X(T, \lambda_i). \quad (3.14)$$

The constraint of  $\mathcal{L}$  (3.13) is well known and is often called Zakharov reduction when  $n = 1$  [25, 11, 1, 3]. In [3], using the  $\bar{\partial}$  method, the authors demonstrated that (3.13) is a result of symmetry constraint of the dKP hierarchy. Here we have shown that (3.13) can be obtained by the dispersionless limit of (2.6), i.e., the dispersionless limit of symmetry constraint of the Sato operator for the dispersionfull hierarchy leads to the symmetry constraint of the Sato function for the dispersionless hierarchy. From (3.3), (3.12), (3.14) and by a tedious computation, we obtain the following equations of hydrodynamical type

$$p_{i,T_k} = [\mathcal{B}_k(p)|_{p=p_i}]_X, \quad (3.15a)$$

$$v_{i,T_k} = [v_i\left(\frac{\partial}{\partial p}\mathcal{B}_k(p)\right)|_{p=p_i}]_X, \quad i = 1, \dots, N. \quad (3.15b)$$

Taking the dispersionless limit of (3.4), we will get the constrained dKP hierarchy reduced by (3.13)

$$(\mathcal{L}^n)_{T_k} = \{\mathcal{B}_k, \mathcal{L}^n\}, \quad (3.16a)$$

$$p_{i,T_k} = [\mathcal{B}_k(p)|_{p=p_i}]_X, \quad (3.16b)$$

$$v_{i,T_k} = [v_i\left(\frac{\partial}{\partial p}\mathcal{B}_k(p)\right)|_{p=p_i}]_X, \quad i = 1, \dots, N. \quad (3.16c)$$

Adding the term  $(\mathcal{B}_k)_{T_n}$  to the right hand side of (3.16a), or taking the dispersionless limit of (2.9) directly, we will obtain the dKP hierarchy with self-consistent sources as

$$(\mathcal{B}_k)_{T_n} - (\mathcal{L}^n)_{T_k} + \{\mathcal{B}_k, \mathcal{L}^n\} = 0, \quad (3.17a)$$

$$p_{i,T_k} = [\mathcal{B}_k(p)|_{p=p_i}]_X, \quad (3.17b)$$

$$v_{i,T_k} = [v_i\left(\frac{\partial}{\partial p}\mathcal{B}_k(p)\right)|_{p=p_i}]_X, \quad i = 1, \dots, N. \quad (3.17c)$$

If " $T_n$ " is viewed as the evolution variable, (3.16) may be regarded as the stationary system of (3.17) like the dispersionfull case. It is not difficult to prove that under (3.17b)

and (3.17c), the equation (3.17a) will be obtained by the compatibility of the following conservation equations

$$p_{T_k} = (\mathcal{B}_k(p))_X, \quad (3.18a)$$

$$p_{T_n} = (\mathcal{L}^n(p))_X = [\mathcal{B}_n(p) + \sum_{i=1}^N \frac{v_i}{p - p_i}]_X. \quad (3.18b)$$

For example, when  $n = 3$ ,  $k = 2$ ,

$$\mathcal{L}^3 = p^3 + 3U_1p + 3U_2 + \sum_{i=1}^N \frac{v_i}{p - p_i}, \quad \mathcal{B}_2 = p^2 + 2U_1,$$

(3.17) becomes the dKP equation with self-consistent sources ( $U_2$  is eliminated by  $U_{2,X} = \frac{1}{2}U_{1,Y}$  and  $T = T_3$ ,  $Y = T_2$ )

$$(U_{1,T} - 3U_1U_{1,X} + \sum_{i=1}^N v_{i,X})_X = \frac{3}{4}U_{1,YY}, \quad (3.19a)$$

$$p_{i,Y} = (p_i^2 + 2U_1)_X, \quad (3.19b)$$

$$v_{i,Y} = 2(v_i p_i)_X, \quad i = 1, \dots, N. \quad (3.19c)$$

Under (3.19b) and (3.19c), the compatibility of the following conservation equations give rise to (3.19a)

$$p_Y = (p^2 + 2U_1)_X = 2pp_X + 2U_{1,X}, \quad (3.20a)$$

$$\begin{aligned} p_T &= (p^3 + 3U_1p + 3U_2 + \sum_{i=1}^N \frac{v_i}{p - p_i})_X \\ &= 3p^2p_X + 3(U_1p)_X + 3U_{2,X} + \sum_{i=1}^N \frac{v_{i,X}}{p - p_i} - \sum_{i=1}^N \frac{v_i(p_X - p_{i,X})}{(p - p_i)^2}. \end{aligned} \quad (3.20b)$$

where  $U_{2,X} = \frac{1}{2}U_{1,Y}$ .

## 4 Hodograph solutions

Motivated by the dKP case, we would like to solve (3.17) by hodograph transformation provided the conservation equations (3.18). Following [6], one can consider the  $M$ -reductions of (3.18) so that the momentum function  $p$  and  $v_i$ ,  $p_i$ ,  $i = 1, \dots, N$  depend only on a set of functions  $W = (W_1, \dots, W_M)$  with  $W_1 = U_1$  and  $(W_1, \dots, W_M)$  satisfy commuting flows

$$\frac{\partial W}{\partial T_n} = A_n(W) \frac{\partial W}{\partial X}, \quad n \geq 2 \quad (4.1)$$

where the  $M \times M$  matrices  $A_n$  are functions of  $(W_1, \dots, W_M)$  only. In the following, we shall take the dKP equation with self-consistent sources (3.19) for example and show its

solutions in the cases of  $M = 1$  and  $M = 2$ .

1.  $M = 1$

In this case,  $p = p(U_1)$ ,  $v_i = v_i(U_1)$ ,  $p_i = p_i(U_1)$  and (4.1) becomes

$$U_{1,Y} = A(U_1)U_{1,X}, \quad U_{1,T} = B(U_1)U_{1,X}. \quad (4.2)$$

From (3.19b), (3.19c) and (3.20), we will get the following relations respectively

$$\frac{dp_i}{dU_1} \left( \frac{A(U_1)}{2} - p_i \right) = 1, \quad (4.3a)$$

$$\frac{dv_i}{dU_1} \left( \frac{A(U_1)}{2} - p_i \right) = v_i \frac{dp_i}{dU_1}, \quad (4.3b)$$

$$\frac{dp}{dU_1} \left( \frac{A(U_1)}{2} - p \right) = 1, \quad (4.3c)$$

$$\frac{dp}{dU_1} B(U_1) = 3p^2 \frac{dp}{dU_1} + 3p + 3U_1 \frac{dp}{dU_1} + 3 \frac{dU_2}{dU_1} + \sum_{i=1}^N \frac{\frac{dv_i}{dU_1}}{p - p_i} - \sum_{i=1}^N v_i \frac{(\frac{dp}{dU_1} - \frac{dp_i}{dU_1})}{(p - p_i)^2}, \quad (4.3d)$$

which implies

$$B = 3U_1 + \frac{3}{4}A^2 - \sum_{i=1}^N \frac{dv_i}{dU_1}, \quad A = 2 \frac{dU_2}{dU_1}. \quad (4.4)$$

It is easy to verify that with (4.3) and (4.4), (4.2) are compatible. Making the hodograph transformations with the change of variables  $(X, Y, T) \rightarrow (U_1, Y, T)$  and  $X = X(U_1, Y, T)$ , we will get the following hodograph equations for  $X$ ,

$$\frac{\partial X}{\partial Y} = -A, \quad \frac{\partial X}{\partial T} = -B = -3U_1 - \frac{3}{4}A^2 + \sum_{i=1}^N \frac{dv_i}{dU_1}, \quad (4.5)$$

which can be easily integrated as follows

$$X + A(U_1)Y + (3U_1 + \frac{3}{4}A(U_1)^2 - \sum_{i=1}^N \frac{dv_i}{dU_1})T = F(U_1), \quad (4.6)$$

where  $F(U_1)$  is an arbitrary function of  $U_1$ .

If we choose  $A(U_1) = C_0 = \text{const}$ ,  $F(U_1) = \beta U_1$ , from (4.6), (4.3a) and (4.3b), we get an implicit solution as

$$X + C_0 Y + [3U_1 + \frac{3}{4}C_0^2 - \sum_{i=1}^N c_i (\frac{C_0^2}{4} - 2(U_1 + d_i))^{-\frac{3}{2}}]T = \beta U_1, \quad (4.7a)$$

$$v_i = c_i [\frac{C_0^2}{4} - 2(U_1 + d_i)]^{-\frac{1}{2}}, \quad (4.7b)$$

$$p_i = \frac{C_0}{2} \pm \sqrt{\frac{C_0^2}{4} - 2(U_1 + d_i)}, \quad i = 1, \dots, N, \quad (4.7c)$$

where  $c_i, d_i, i = 1, \dots, N$  are constants. If  $d_1 = d_2 = \dots = d_N$  and  $\sum_{i=1}^N c_i = 0$ , (4.7a) degenerates to the solution of dKP equation [6].

If we assume  $v_1 = v_2 = \dots = v_N$  and  $\sum_{i=1}^N \frac{dv_i}{dU_1} = 3U_1$ , i.e.,  $\frac{dv_i}{dU_1} = \frac{3}{N}U_1$ , and taking  $F(U_1) = 0$ , by a direct computation, we will obtain an explicit solution of the dKP equation with self-consistent sources (3.19) as follows

$$U_1 = \frac{4Y^2 - 6TX \pm 4Y\sqrt{Y^2 - 3TX}}{225T^2},$$

$$v_i = \frac{3}{2N}U_1^2,$$

$$p_i = 2\sqrt{2U_1}, \quad i = 1, \dots, N.$$

2.  $M = 2$

In this case we denote  $W_1 = U_1, W_2 = W$ , then  $v_i = V_i(U_1, W), p_i = p_i(U_1, W)$  and  $p = p(U_1, W)$  with the commuting flows

$$\begin{pmatrix} U_1 \\ W \end{pmatrix}_Y = A \begin{pmatrix} U_1 \\ W \end{pmatrix}_X, \quad \begin{pmatrix} U_1 \\ W \end{pmatrix}_T = B \begin{pmatrix} U_1 \\ W \end{pmatrix}_X, \tag{4.8}$$

where  $A = (A_{ij})$  and  $B = B_{ij}$  are  $2 \times 2$  matrix functions of  $U_1$  and  $W$ . Requiring  $U_{1,x}$  and  $W_x$  are independent, (3.19b), (3.19c) and (3.20) give rise to the following relations respectively

$$\left(\frac{\partial p_i}{\partial U_1}, \frac{\partial p_i}{\partial W}\right)A = \left(2 + 2p_i \frac{\partial p_i}{\partial U_1}, 2p_i \frac{\partial p_i}{\partial W}\right), \tag{4.9a}$$

$$\left(\frac{\partial v_i}{\partial U_1}, \frac{\partial v_i}{\partial W}\right)A = \left(2 \frac{\partial(v_i p_i)}{\partial U_1}, 2 \frac{\partial(v_i p_i)}{\partial W}\right), \tag{4.9b}$$

$$\left(\frac{\partial p}{\partial U_1}, \frac{\partial p}{\partial W}\right)A = 2p \left(\frac{\partial p}{\partial U_1}, \frac{\partial p}{\partial W}\right) + (2, 0), \tag{4.9c}$$

$$\begin{aligned} \left(\frac{\partial p}{\partial U_1}, \frac{\partial p}{\partial W}\right)B &= 3p^2 \left(\frac{\partial p}{\partial U_1}, \frac{\partial p}{\partial W}\right) + 3 \left(\frac{\partial(U_1 p)}{\partial U_1}, \frac{\partial(U_1 p)}{\partial W}\right) + 3 \left(\frac{\partial U_2}{\partial U_1}, \frac{\partial U_2}{\partial W}\right) \\ &+ \sum_{i=1}^N \frac{1}{p-p_i} \left(\frac{\partial v_i}{\partial U_1}, \frac{\partial v_i}{\partial W}\right) - \sum_{i=1}^N \frac{v_i}{(p-p_i)^2} \left(\frac{\partial p}{\partial U_1} - \frac{\partial p_i}{\partial U_1}, \frac{\partial p}{\partial W} - \frac{\partial p_i}{\partial W}\right), \end{aligned} \tag{4.9d}$$

which implies  $A(U_1, W)$  and  $B(U_1, W)$  must satisfy

$$B = \frac{3}{4}A^2 + 3U_1 I - \sum_{i=1}^N \frac{\partial v_i}{\partial U_1} I - \begin{pmatrix} 0 & \sum_{i=1}^N \frac{\partial v_i}{\partial W} \\ \frac{A_{11}}{A_{12}} \sum_{i=1}^N \frac{\partial v_i}{\partial W} & \frac{A_{22}-A_{11}}{A_{12}} \sum_{i=1}^N \frac{\partial v_i}{\partial W} \end{pmatrix} \tag{4.10}$$

with  $A_{11} = 2 \frac{\partial U_2}{\partial U_1}$  and  $A_{12} = 2 \frac{\partial U_2}{\partial W}$ .

For simplicity we assume  $\frac{\partial v_i}{\partial W} = 0, i = 1, \dots, N$ . Then

$$B = \frac{3}{4}A^2 + 3U_1 I - \sum_{i=1}^N \frac{\partial v_i}{\partial U_1} I = \frac{3}{4}(tr A)A + 3(U_1 - \frac{1}{4}det A - \frac{1}{3} \sum_{i=1}^N \frac{\partial v_i}{\partial U_1})I, \tag{4.11}$$

where the formula  $A^2 = (\text{tr}A)A - (\det A)I$  is used.

With (4.11), the compatibility for (4.8) requires  $A$  to satisfy

$$\left( \frac{\frac{1}{4} \frac{\partial \det A}{\partial W}}{\frac{\partial(U_1 - \frac{1}{4} \det A - \frac{1}{3} \sum_{i=1}^N \frac{\partial v_i}{\partial U_1})}{\partial U_1}} \right) = A \left( \frac{\frac{1}{4} \frac{\partial \text{tr} A}{\partial W}}{-\frac{1}{4} \frac{\partial \text{tr} A}{\partial U_1}} \right). \quad (4.12)$$

Using the hodograph transformation changing the independent variables  $(X, Y, T)$  to  $(U_1, W, T)$  with  $X = X(U_1, W, T)$  and  $Y = Y(U_1, W, T)$ , we get

$$\begin{pmatrix} -X_W \\ X_{U_1} \end{pmatrix} = A \begin{pmatrix} Y_W \\ -Y_{U_1} \end{pmatrix}, \quad \begin{pmatrix} \frac{\partial(X, Y)}{\partial(W, T)} \\ -\frac{\partial(X, Y)}{\partial(U_1, T)} \end{pmatrix} = B \begin{pmatrix} Y_W \\ -Y_{U_1} \end{pmatrix}, \quad (4.13)$$

where  $\frac{\partial(X, Y)}{\partial(W, T)} = X_W Y_T - X_T Y_W$ . It is not difficult to see that (4.13) has solutions in the form

$$X + 3(U_1 - \frac{1}{4} \det A - \frac{1}{3} \sum_{i=1}^N \frac{\partial v_i}{\partial U_1})T = F(U_1, W), \quad (4.14a)$$

$$Y + \frac{3}{4}(\text{tr}A)T = G(U_1, W), \quad (4.14b)$$

where we have required that  $Y_{U_1}$  and  $Y_W$  are independent, while  $F$  and  $G$  satisfy the linear equations

$$\begin{pmatrix} -F_W \\ F_{U_1} \end{pmatrix} = A \begin{pmatrix} G_W \\ -G_{U_1} \end{pmatrix}. \quad (4.15)$$

An example of solution is given by

$$A = \begin{pmatrix} W & U_1 \\ 4 & W \end{pmatrix}, \quad (4.16)$$

and

$$v_i = c_i U_1, \quad p_i = \frac{W}{2}, \quad i = 1, \dots, N, \quad (4.17)$$

with  $c_i, i = 1, \dots, N$  are constants.

(4.14) now becomes

$$X + 3[U_1 - \frac{1}{4}(W^2 - 4U_1) - \frac{1}{3} \sum_{i=1}^N c_i]T = F(U_1, W), \quad (4.18a)$$

$$Y + \frac{3}{4}2WT = G(U_1, W). \quad (4.18b)$$

From (4.15) and  $F_{WU_1} = F_{U_1W}$ ,  $G$  must satisfy

$$2G_{U_1} + U_1 G_{U_1 U_1} - 4G_{WW} = 0.$$

Taking  $G = -W$  and from (4.15),  $F = \frac{1}{2}W^2 - 4U_1$  and we obtain a solution of (3.19) as follows

$$U_1 = \frac{Y^2}{2(3T+2)^2} + \frac{(\sum_{i=1}^N c_i)T - X}{2(3T+2)}, \quad (4.19a)$$

$$v_i = c_i U_1 = c_i \left[ \frac{Y^2}{2(3T+2)^2} + \frac{(\sum_{i=1}^N c_i)T - X}{2(3T+2)} \right], \quad (4.19b)$$

$$p_i = \frac{W}{2} = -\frac{Y}{3T+2}, \quad i = 1, \dots, N. \quad (4.19c)$$

(4.19) is a global solution for  $T > -\frac{2}{3}$  and (4.19a) degenerates to the solution of the dKP equation [6] when  $\sum_{i=1}^N c_i = 0$ .

**Acknowledgments.** *This work was supported by the Chinese Basic Research Project “Nonlinear Science”.*

## References

- [1] AOYAMA S and KODAMA Y, Topological Landau-Ginzburg theory with a rational potential and the dispersionless KP hierarchy, *Commun. Math. Phys.* **182** (1996), 185–219.
- [2] BOGDANOV L V, KONOPELCHENKO B G and MARTINEZ ALONSO L, Quasiclassical D-bar method: Generating equations for dispersionless integrable hierarchies, preprint [arXiv:nlin.SI/0111062](https://arxiv.org/abs/nlin.SI/0111062).
- [3] BOGDANOV L V and KONOPELCHENKO B G, Symmetry constraints for the dispersionless integrable equations and systems of hydrodynamical type, preprint [arXiv:nlin.SI/0312013](https://arxiv.org/abs/nlin.SI/0312013).
- [4] CHENG Y, Modifying the KP, the nth Constrained KP Hierarchies and their Hamiltonian Structures, *Commun. Math. Phys.* **171** (1995), 661–682.
- [5] DICKEY L A, Soliton equation and Hamiltonian systems, World Scientific, Singapore, 1991.
- [6] KODAMA Y, A method for solving the dispersionless KP equation and its exact solutions, *Phys. Lett. A* **129** (1988), 223–226; Solutions of the dispersionless Toda equation, **147** (1990), 477–482.
- [7] KODAMA Y and GIBBONS J, A method for solving the dispersionless KP hierarchy and its exact solutions II, *Phys. Lett. A* **135** II (1989), 167–170.
- [8] KONOPELCHENKO B, MARTINEZ ALONSO L and RAGNISCO O, The D-bar Approach to the dispersionless KP hierarchy, *J. Phys. A* **34** (2001), 10209–10217.
- [9] KONOPELCHENKO B, MARTINEZ ALONSO L and MEDINA E, Quasiconformal mappings and solutions of the dispersionless KP hierarchy, preprint [arXiv:nlin.SI/0202013](https://arxiv.org/abs/nlin.SI/0202013).
- [10] KRICHEVER I M, The averaging method for two-dimensional “integrable” equations, *Func. Anal. Priloz.* **22** (1988), 37–52.
- [11] KRICHEVER I M, Topological minimal models and dispersionless Lax equations, *Commun. Math. Phys.* **143** (1992), 415–429.

- 
- [12] KUPERSCHMIDT B A and MANIN YU I, Long wave equations with a free surface. I. Conservation laws and solutions, *Func. Anal. Appl.* **11** (1977) [in Russian], 31–42; Long wave equations with a free surface. II. The Hamiltonian structure and the higher equations, *Func. Anal. Appl.* **17** (1978), 25–37.
- [13] LAX P D and LEVERMORE C D, The small dispersion limit of the Korteweg-de Vries equation I *Commun. Pure Appl. Math.* **36** (1983) 253–290; The small dispersion limit of the Korteweg-de Vries equation II, *Commun. Pure Appl. Math.* **36** (1983) 571–593; The small dispersion limit of the Korteweg-de Vries equation III, *Commun. Pure Appl. Math.* **36** (1983) 809–830.
- [14] LEBEDEV D and MANIN YU I, Conservation Laws and Lax Representation on Benney’s Long Wave Equations, *Phys. Lett. A* **74** (1979), 154–156.
- [15] LEON J and LATIFI A, Solutions of an initial-boundary value problem for coupled non-linear waves, *J. Phys. A* **23** (1990), 1385–1403.
- [16] MARTINEZ ALONSO L and MANÑAS M, Additional symmetries and solutions of the dispersionless KP hierarchy, preprint [arXiv:nlin.SI/0302009](https://arxiv.org/abs/nlin.SI/0302009).
- [17] MEL’NIKOV V K, A direct method for deriving a multi-soliton solution for the problem of interaction of waves on the  $x, y$  plane, *Commun. Math. Phys.* **112** (1987), 639–652.
- [18] MEL’NIKOV V K, Interaction of solitary waves in the system described by the Kadomtsev-Petviashvili equation with a self-consistent source, *Commun. Math. Phys.* **126** (1989), 201–215.
- [19] OEVEL W and STRAMPP W, Constrained KP Hierarchy and Bi-Hamiltonian Structures, *Commun. Math. Phys.* **157** (1993), 51–81.
- [20] SATO M, Soliton equations as dynamical systems on infinite dimensional Grassmann manifolds, *RIMS Kokyuroku (Kyoto Univ.)* **439** (1980), 30.
- [21] TAKASAKI K and TAKEBE T, SDIFF(2) KP hierarchy, *Int. J. Mod. Phys. A Suppl.* **1B** (1992), 889–922.
- [22] TAKASAKI K and TAKEBE T, Integrable hierarchies and dispersionless limit, *Rev. Math. Phys.* **7** (1995), 743–808.
- [23] XIAO T and ZENG Y, Generalized Darboux transformations for the KP equation with self-consistent sources, *J. Phys. A* **37** (2004), 7143–7162.
- [24] XIAO T and ZENG Y, A new constrained mKP hierarchy and the generalized Darboux transformation for the mKP equation with self-consistent sources, *Physica A* **353** (2005), 38–60.
- [25] ZAKHAROV V E, Benney equation and quasiclassical approximation in the inverse problem method, *Func. Anal. Priloz.* **14** (1980) 89–98; On the Benney equation, *Physica D* **3** (1981), 193–202.
- [26] ZAKHAROV V E, Dispersionless limit of integrable systems in (2+1)-dimensions, in *Singular limit of dispersive waves*, Editors: Erconali N M *et al*, Plenum, New York, 1994, 165–174.
- [27] ZENG Y, MA W and LIN R, Integration of the soliton hierarchy with self-consistent sources, *J. Math. Phys.* **41** (2000), 8.
- [28] ZENG Y, SHAO Y and MA W, Integral-type Darboux transformations for the mKdV hierarchy with self-consistent sources, *Commun. Theor. Phys.* **38** (2002), 641–648.