



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantis-pess.com/journals/artres>

P152: MICROALBUMINURIA IN NEWLY DIAGNOSED DIABETES MELLITUS: NOT ONLY ABOUT BLOOD PRESSURE OR ARTERIAL STIFFNESS

Barbara Nikolaidou, Eleni Gavriilaki, Eugene Gkaliagkousi, Ioanna Zografou, Areti Triantafyllou, Antonios Lazaridis, Stella Douma

To cite this article: Barbara Nikolaidou, Eleni Gavriilaki, Eugene Gkaliagkousi, Ioanna Zografou, Areti Triantafyllou, Antonios Lazaridis, Stella Douma (2017) P152: MICROALBUMINURIA IN NEWLY DIAGNOSED DIABETES MELLITUS: NOT ONLY ABOUT BLOOD PRESSURE OR ARTERIAL STIFFNESS, Artery Research 20:C, 102–103, DOI: <https://doi.org/10.1016/j.artres.2017.10.173>

To link to this article: <https://doi.org/10.1016/j.artres.2017.10.173>

Published online: 7 December 2019

Japan (cardio-ankle vascular index) method. Blood samples were taken for ELISA assays. Analysis was completed with SPSS software

Results: Forty patients were evaluated in this preliminary project (29 male/11 female, mean age 70.7 ± 11.99), with four experiencing a further event during the six month follow up (10%).

All biomarkers and both measurements for arterial stiffness had a higher mean value in patients with a further event (hsCRP 3.89 vs 1.42, $P = 0.08$; EPO 9.06 vs 9.01, $P = 0.85$; sRANKL 0.05 vs 0.03, $P = 0.31$; PRDX1 6.27 vs 6.21, $P = 0.95$; CAVI 11.13 vs 9.69, $P = 0.15$; cfPWV 10.82 vs 10.2, $P = 0.55$), however none were statistically significant.

Levels of PRDX1 were elevated acutely post-event before falling significantly ($R = -0.475$, $P = 0.002$), while hsCRP and EPO continued to be elevated at >10 days post-event.

In addition, CAVI correlated closely with hsCRP ($R = 0.28$, $P = 0.09$) and EPO ($R = 0.29$, $P = 0.08$), but cfPWV was not closely related to any of the biomarkers.

Conclusions: This preliminary data suggests that biomarkers, particularly EPO and hsCRP, are more closely related to CAVI than cfPWV. hsCRP was the most relevant as an independent predictive factor for further vascular events.

Poster Session II – Obesity and Diabetes

P150

ARTERIAL STIFFNESS AND PROGRESSION OF CEREBRAL WHITE MATTER LESIONS IN ASYMPTOMATIC PATIENTS WITH TYPE 2 DIABETES AND MATCHED CONTROLS: A 5-YEAR COHORT STUDY

Kristian Løkke Funck¹, Esben Laugesen¹, Pernille Høyem¹, Brian Stausbøll-Grøn², Won Kim^{3,2}, Leif Østergaard⁴, Dora Zeidler⁴, Troels Krarup Hansen¹, Per Løgstrup Poulsen¹

¹Department of Endocrinology and Internal Medicine, Aarhus University Hospital, Denmark

²MR Research Centre, Department of Clinical Medicine, Aarhus University, Denmark

³Department of Cardiology, Aarhus University Hospital, Denmark

⁴Center of Functionally Integrative Neuroscience, Aarhus University Hospital, Denmark

Aim: Stroke is a frequent and feared complication in patients with type 2 diabetes. Arterial stiffness may improve current suboptimal risk prediction of stroke. However, studies in diabetes populations are lacking. We investigated the association between arterial stiffness progression (carotid-femoral pulse wave velocity [PWV]) and the progression of cerebral white matter lesions (WML), a marker of stroke risk, in patients with type 2 diabetes and matched controls.

Methods: In a 5-year follow-up study, data from 49 patients and 58 controls were available for analysis. At baseline, participants had a mean (\pm SD) age of 59 ± 10 years and patients had a median (range) diabetes duration of 1.8 (1.0–3.2) years. Fifty-two (49%) were males. At both baseline and follow-up, PWV was obtained by tonometry and WML by cerebral T2-FLAIR MRI. WML was assessed by Breteler score, and progression was defined as an upward change in category during follow-up.

Results: Patients with type 2 diabetes had a higher PWV than controls at both baseline (9.2 ± 2.2 vs. 7.9 ± 1.4 m/s, $p < 0.01$) and follow-up (9.8 ± 2.4 vs. 8.6 ± 1.9 m/s, $p = 0.01$). Breteler scores and WML progression were similar in the two groups ($p > 0.05$). PWV progression was associated with WML progression in the total cohort (adjusted for age, sex, diabetes, baseline PWV and systolic blood pressure progression: OR 1.58 [95%CI: 1.09–2.28], $p = 0.02$). We found no interaction between diabetes and PWV progression on WML progression.

Conclusions: PWV progression is associated with WML progression in patients with type 2 diabetes and healthy controls. PWV candidates as a new risk marker for stroke.

P151

CENTRAL PULSE PRESSURE IS ASSOCIATED WITH AORTIC-BRACHIAL STIFFNESS MISMATCH IN PATIENTS WITH ARTERIAL HYPERTENSION AND DIABETES MELLITUS

Elena Troitskaya, Ekaterina Starostina, Zhanna Kobalava
RUDN University, Russia

Background: Central pulse pressure (PP) is a surrogate measure of arterial stiffness (AS) and a predictor of cardiovascular events in type 2 diabetes mellitus (T2DM). AS gradient reflects the vascular ageing.

The aim of the study: To evaluate the associations between 24-h central PP and parameters of AS in patients with arterial hypertension (AH) and T2DM. **Materials and methods:** 90 patients with AH and T2DM were included (39% males, mean age 63.8 ± 11.6). Mean office BP was $146 \pm 23/86 \pm 10$ mmHg. Median duration of DM was 8.5 years. 24-h peripheral and central BP monitoring was performed (BPLab Vasotens). AS parameters were assessed by applanation tonometry (Sphygmocor, AtCor). AS gradient was calculated as cfPWV/crPWV and its elevation ≥ 1 was considered as AS mismatch. $p < 0.05$ was considered significant.

Results: Mean central BP was $132 \pm 18/79 \pm 12$ mmHg, cfPWV -10.5 ± 2.4 m/s, crPWV -8.5 ± 1.4 m/s, AS gradient -1.2 ± 0.3 . 24-h central BP levels were as follows: $132 \pm 18/79 \pm 12$ mmHg for 24hBP, $132 \pm 18/81 \pm 13$ mmHg for daytime and $129 \pm 20/75 \pm 11$ mmHg for nighttime. Mean PP levels were 53 ± 14 , 52 ± 14 and 56 ± 15 mmHg, respectively. 24-h central PP increase > 50 mmHg was observed in 30% patients. These patients were older (67.1 ± 10.5 vs 61.8 ± 11.9 years), had higher median duration of DM (10; IQR 7–15 vs 5; IQR 0, 9–12 years), cfPWV (11.4 ± 1.9 vs 10.0 ± 2.6 m/s) and stiffness gradient (1.4 ± 0.3 vs 1.2 ± 0.2), $p < 0.05$ for trend. There were significant correlations between 24-h central PP and age ($r = 0.27$) and AS gradient ($r = 0.32$), $p < 0.05$ for trend. No predictors of PP elevation were found.

Conclusions: In diabetic patients with AH increase of central PP is associated with aortic-brachial stiffness mismatch. This finding confirms its importance as a marker of vascular ageing in this patient population.

P152

MICROALBUMINURIA IN NEWLY DIAGNOSED DIABETES MELLITUS: NOT ONLY ABOUT BLOOD PRESSURE OR ARTERIAL STIFFNESS

Barbara Nikolaidou¹, Eleni Gavriilaki¹, Eugene Kkaliagkousi¹, Ioanna Zografou², Areti Triantafyllou¹, Antonios Lazaridis¹, Stella Douma¹

¹Third Department of Internal Medicine, Papageorgiou General Hospital of Thessaloniki, Greece

²Hippokraton General Hospital of Thessaloniki, Greece

Background: Diabetes mellitus (DM) and hypertension (EH) are both associated with micro- and macro-vascular damage. Microalbuminuria is a recognized marker of sub-clinical target organ damage in both DM and EH. However, it's determinants in newly diagnosed DM with or without EH remain unclear.

Methods: We enrolled consecutive newly diagnosed DM patients, recording history, demographics, renal, lipid and glycemic profile, office and ambulatory blood pressure, macro-(pulse-wave velocity/PWV) and micro-vascular (microalbuminuria in 24-hour urine) damage and subclinical atherosclerosis (intima-media thickness).

Results: We studied 65 DM patients (40 male: 25 female, aged 57 ± 11 years), with a median duration from diagnosis of 2 weeks. Their fasting glucose was 121.5 (IR: 36) mg/dl, HbA1c: 7.47 (IR: 2%). Among them, 26 had already been diagnosed with EH (median duration of 8 (IR: 8) years), while 17 were diagnosed with EH at the time of DM diagnosis. No difference was observed between the two groups, except for significantly higher office and ambulatory BP and PWV in the newly diagnosed EH patients. Microalbuminuria was associated with fasting glucose ($p = 0.04$), HbA1c ($p = 0.002$), serum creatinine ($p = 0.035$), glomerular filtration rate (GFR) ($p = 0.002$), office systolic ($p = 0.009$) and diastolic ($p = 0.026$) BP and PWV ($p = 0.031$). In the multivariate analysis, HbA1c (beta = 0.351, $p = 0.015$) was the only determinant of microalbuminuria.

Conclusions: Our study indicates that hyperglycaemia has a significant impact on microalbuminuria even in patients with newly diagnosed DM and

EH (either newly diagnosed with high BP values or longer lasting), emphasizing on the need of early and effective glycemetic control.

P153

MARKER OF TYPE VI COLLAGEN FORMATION (PRO-C6) IS ASSOCIATED WITH HIGHER ARTERIAL STIFFNESS IN TYPE 1 DIABETES

Tine Willum Hansen¹, Daniel G. K. Rasmussen², Simone Theilade¹, Signe H. Nielsen², Morten A. Karsdal², Federica Genovese², Peter Rossing¹

¹Steno Diabetes Center Copenhagen, Gentofte, Denmark

²Nordic Bioscience, Herlev, Denmark

Aim: Arterial stiffening reflects fragmentation and loss of elastin fibers and accumulation of stiffer collagen fibers in the media of large arteries. We evaluated associations between carotid-femoral pulse wave velocity (cfPWV) and a marker of collagen type VI formation (Pro-C6) and a marker of collagen type III degradation (C3M) in type 1 diabetes.

Methods: Serum and urinary level of Pro-C6 and C3M was measured with ELISA in 634 patients with type 1 diabetes. cfPWV was assessed by the SphygmoCor device. We applied unadjusted and adjusted linear regression analyses. Adjustment included sex, age, mean arterial pressure, LDL cholesterol, smoking, HbA_{1c}, eGFR and urinary albumin excretion rate. To adjust for urine output levels, the urinary markers were normalized for urinary creatinine.

Results: Of the 634 patients, 349 (55%) were male, mean \pm SD age was 54.6 ± 12.6 years and cfPWV 10.4 ± 3.3 m/s.

Higher serum and urinary level of Pro-C6 was associated with higher cfPWV in unadjusted models ($p \leq 0.039$), after adjustment only higher serum level remained significantly associated with higher cfPWV (β estimate per doubling: 0.47 ± 0.21 ; $p = 0.028$).

Lower urinary level of C3M was associated with higher cfPWV in the unadjusted model ($p < 0.001$), but significance was lost after adjustment ($p = 0.44$). Higher serum level of C3M was associated with higher cfPWV in the unadjusted model ($p = 0.002$), but significance was lost after adjustment ($p = 0.34$).

Conclusion: In type 1 diabetes, higher serum levels of Pro-C6, a marker of collagen type VI formation, was associated with increased arterial stiffness. This observation could introduce a new target for therapeutic intervention.

P154

VASCULAR STRUCTURE AND FUNCTION IN RELATION TO WEIGHT EXCESS AND BLOOD PRESSURE IN A SAMPLE OF OBESE CHILDREN

Sara Bonafini¹, Angela Tagetti¹, Francesco Dalle Vedove¹, Sara Raimondi¹, Alice Giontella¹, Paolo Cavarzere², Rossella Gaudino², Pietro Minuz¹, Claudio Maffei², Franco Antoniazzi², Cristiano Fava¹

¹University of Verona, Department of Medicine, Italy

²University of Verona, Department of Surgery, Dentistry, Paediatrics and Gynaecology, Italy

Objective: To investigate the influence of weight excess and blood pressure (BP) on vascular structure and function in a sample of obese children.

Methods: We measured office and ambulatory BP (ABPM), carotid intima-media thickness (cIMT), endothelial function by the Flow Mediated Dilation (FMD) technique, carotid distensibility (cDC) (ultrasonograph: LogiQ P5 Pro; software: Multimedia Video Engine II, DSP Lab, Pisa CNR, Italy) (1) and stiffness index (SI) by photoplethysmography (Pulce Trace PT1000, MicroMedical Ltd, UK) (2) in overweight and obese children (BMI > 90th percentile for sex and age).

Results: Seventy children with weight excess were enrolled (age 11.5 ± 2.4 years; female n: 30). cDC showed inverse correlation with BMI and waist circumference ($r_{\text{Spearman}} = -0.403$ and $r_s = -0.346$, respectively), 24h-SBP ($r_s = -0.449$) and nighttime-SBP ($r_s = -0.490$). SI inversely correlated with BMI ($r_s = -0.359$) and waist/height ratio ($r_s = -0.303$). When comparing normotensive and hypertensive children, as defined on the basis of the

ABPM, no significant differences in vascular tests were found, although cDC tended to be lower in hypertensive. In normotensive subjects (n: 53) cIMT directly correlated with nighttime-SBP and DBP. Most of the correlations remained significant after adjustment for age, sex, BMI and BP.

Conclusions: These data suggest that arterial elasticity is negatively affected by weight excess and 24h-BP levels even in childhood. In the normotensive subgroup it is detectable an effect of BP also on arterial structure.

References

(1) Doyon A, et al., Carotid Artery Intima-Media Thickness and Distensibility in Children and Adolescents: Reference Values and Role of Body Dimension. *Hypertension* 2013; 62: 550-556 (2) Millesseu SC et al., Determination of age-related increases in large artery stiffness by digital pulse contour analysis. *Clin Sci (Lond)* 2002; 103:371-7.

P155

CORRELATION OF SOLUBLE RECEPTOR FOR ADVANCED GLYCATION END PRODUCTS AND S100-A1 ON ARTERIAL STIFFNESS IN NORMOTENSIVE PATIENTS WITH DIABETES

Patricia Quezada Fernandez^{1,2}, Fernando Grover Páez^{3,4,5}, Becerra Ramos^{6,7}, Carlos^{6,7}, Jhonatan Trujillo Quiros⁶, Walter Trujillo Rangel⁶, Sánchez Rodríguez Sánchez Rodríguez⁶, David Cardona Müller^{6,8}, Mayra Jimenez Cázarez⁶

¹Experimental Therapeutic and Clinic Institute, Physiology Department of the University of Guadalajara, Mexico

²PNPC, National Council of Science and Technology (CONACYT), Mexico

³University of Guadalajara, Physiology Department of the University of Guadalajara, Mexico

⁴Experimental Therapeutic and Clinic Institute, University of Guadalajara, Mexico

⁵National System of Researchers Grade 1, National Council of Science and Technology, Mexico

⁶Vascular Mechanics Laboratory, Experimental Therapeutic and Clinic Institute, Physiology Department, Health Sciences University Center, University of Guadalajara, Mexico

⁷National System of Researchers, National Council of Science and Technology (CONACYT), Mexico

⁸National Council of Cardiology, Mexico

Background: Accumulation of advanced glycation end products (AGEs) are involved in several pathophysiological processes in the vessel wall, that may cause premature atherosclerosis and arterial stiffening (1). A soluble form of RAGE (sRAGE), which is a splice variant of full-length RAGE has been considered to be protective against diseases originating from RAGE activation since sRAGE can bind and sequester RAGE ligands and reduce RAGE activation (2). S100A1 is the most abundant calcium-binding protein in myocardial tissue and is a major determinant of cardiac function. This circulating ligand of the RAGE is known as a pro-inflammatory factor in diabetes. Aberrant expression levels of S100A1 surfaced as molecular key defects, driving the pathogenesis of cardiovascular diseases (3).

Objective: This study was design to explore the relationship between serum levels of sRAGE and S100A1 on arterial stiffness in non-hypertensive patients with diabetes.

Methods: Using a cross-sectional design, a total of 20 non-hypertensive patients with diabetes were recruited. A fasting blood sample, medical history and arterial stiffness parameters were collected.

Results: In bivariate analysis, sRAGE positively correlated with time evolution of diabetes ($r = 0.503$, $p < 0.024$) and negatively correlated with systolic ($r = -0.457$, $p = 0.043$) and diastolic blood pressure ($r = -0.527$, $p = 0.017$). S100A1 positively correlated with creatinine ($r = 0.724$, $p < 0.000$) and negatively correlated with peripheral and central hemodynamics, including augmentation index ($r = -0.469$, $p = 0.037$), as shown in Table 2.