



Journal of Nonlinear Mathematical Physics

ISSN (Online): 1776-0852

ISSN (Print): 1402-9251

Journal Home Page: <https://www.atlantis-press.com/journals/jnmp>

Multicomplex solitons

Julia Cen, Andreas Fring

To cite this article: Julia Cen, Andreas Fring (2020) Multicomplex solitons, Journal of Nonlinear Mathematical Physics 27:1, 17–35, DOI: <https://doi.org/10.1080/14029251.2020.1683963>

To link to this article: <https://doi.org/10.1080/14029251.2020.1683963>

Published online: 04 January 2021

Multicomplex solitons

Julia Cen

*Department of Mathematics, City, University of London
Northampton Square, London EC1V 0HB, UK
julia.cen.1@city.ac.uk*

Andreas Fring

*Department of Mathematics, City, University of London
Northampton Square, London EC1V 0HB, UK
a.fring@city.ac.uk*

Received 7 December 2018

Accepted 11 June 2019

We discuss integrable extensions of real nonlinear wave equations with multi-soliton solutions, to their bicomplex, quaternionic, coquaternionic and octonionic versions. In particular, we investigate these variants for the local and nonlocal Korteweg-de Vries equation and elaborate on how multi-soliton solutions with various types of novel qualitative behaviour can be constructed. Corresponding to the different multicomplex units in these extensions, real, hyperbolic or imaginary, the wave equations and their solutions exhibit multiple versions of antilinear or \mathcal{PT} -symmetries. Utilizing these symmetries forces certain components of the conserved quantities to vanish, so that one may enforce them to be real. We find that symmetrizing the noncommutative equations is equivalent to imposing a \mathcal{PT} -symmetry for a newly defined imaginary unit from combinations of imaginary and hyperbolic units in the canonical representation.

Keywords: nonlinear wave equations; solitons; quaternions; coquaternions; octonions

2000 Mathematics Subject Classification: 34A34, 35Q51, 35Q53, 74J35

1. Introduction

The underlying mathematical structure of quantum mechanics, a Hilbert space over the field of complex numbers, can be generalized and modified in various different ways. One may for instance re-define the inner product of the Hilbert space or alter, typically enlarge, the field over which this space is defined. The first approach has been pursued successfully since around twenty years [8], when it was first realized that the modification of the inner product allows to include non-Hermitian Hamiltonians into the framework of a quantum mechanical theory. When these non-Hermitian Hamiltonians are \mathcal{PT} -symmetric/quasi-Hermitian [7, 28, 32] they possess real eigenvalues when their eigenfunctions are also \mathcal{PT} -symmetric or pairs of complex conjugate eigenvalues when the latter is not the case. The reality of the spectrum might only hold in some domain of the coupling constant, but break down at what is usually referred to as an exceptional point when at least two eigenvalues coalesce. Higher order exceptional points may occur for larger degeneracies. In order to unravel the structure of the neighbourhood of these points one can make use of the second possibility of generalizations of standard quantum mechanics and change the type of fields over which the Hilbert space is defined. This view helps to understand the bifurcation structure at these points and has been recently investigated for the analytically continued Gross-Pitaevskii equation

with bicomplex interaction terms [16, 18, 21]. In a similar spirit, systems with finite dimensional Hilbert spaces have been formulated over Galois fields [37]. Hyperbolic extensions of the complex Hilbert space have been studied in [38]. The standard Schrödinger equation was bicomplexified in [3] and further studied in [4–6, 35]. Quaternionic and coquaternionic quantum mechanics and quantum field theory have been studied for a long time, see e.g. [2, 19, 20], mainly motivated by the fact that they may be related to various groups and algebras that play a central role in physics, such as $SO(3)$, the Lorentz group, the Clifford algebra or the conformal group. Recently it was suggested that they [9] provide a unifying framework for complexified classical and quantum mechanics. Octonionic Hilbert spaces have been utilized for instance in the study of quark structures [22].

Drawing on various relations between the quantum mechanical setting and classical integrable nonlinear systems that possess soliton solutions, such as the formal identification of the L operator in a Lax pair as a Hamiltonian, many of the above possibilities can also be explored in the latter context. Most direct are the analogues of the field extensions. Previously we demonstrated [12, 13] that one may consistently extend real classical integrable nonlinear systems to the complex domain by maintaining the reality of the energy. Here we go further and investigate multicomplex versions of these type of nonlinear equations. We demonstrate how these equations can be solved in several multicomplex settings and study some of the properties of the solutions. We explore three different possibilities to construct solutions that are not available in a real setting, i) using multicomplex shifts in a real solutions, ii) exploiting the complex representations by defining a new imaginary unit in terms of multicomplex ones and iii) exploiting the idempotent representation. We take \mathcal{PT} -symmetry as a guiding principle to select out physically meaningful solutions with real conserved quantities, notably real energies. We clarify the roles played by the different types of \mathcal{PT} -symmetries. For the noncommutative versions, that is quaternionic, coquaternionic and octonionic, we find that imposing certain \mathcal{PT} -symmetries corresponds to symmetrizing the noncommutative terms in the nonlinear differential equations.

Our manuscript is organized as follows: In section 2 we discuss the construction of bicomplex multi-solitons for the standard Korteweg de-Vries (KdV) equation and its nonlocal variant. We present two different types of construction schemes leading to solutions with different types of \mathcal{PT} -symmetries. We demonstrate that the conserved quantities constructed from these solutions, in particular the energy, are real. In section 3, 4 and 5 we discuss solution procedures for noncommutative versions of the KdV equation in quaternionic, coquaternionic and octonionic form, respectively. Our conclusions are stated in section 6.

2. Bicomplex solitons

2.1. Bicomplex numbers and functions

We start by briefly recalling some key properties of bicomplex numbers and functions to settle our notations and conventions. Denoting the field of complex numbers with imaginary unit i as

$$\mathbb{C}(i) = \{x + iy \mid x, y \in \mathbb{R}\}, \quad (2.1)$$

the *bicomplex numbers* \mathbb{B} form an algebra over the complex numbers admitting various equivalent types of representations

$$\mathbb{B} = \{z_1 + Jz_2 \mid z_1, z_2 \in \mathbb{C}(I)\}, \quad (2.2)$$

$$= \{w_1 + Iw_2 \mid w_1, w_2 \in \mathbb{C}(J)\}, \quad (2.3)$$

$$= \{a_1\ell + a_2I + a_3J + a_4k \mid a_1, a_2, a_3, a_4 \in \mathbb{R}\}, \quad (2.4)$$

$$= \{v_1e_1 + v_2e_2 \mid v_1 \in \mathbb{C}(I), v_2 \in \mathbb{C}(J)\}. \quad (2.5)$$

The *canonical basis* is spanned by the units ℓ, I, J, k , involving the two *imaginary units* I and J with $I^2 = J^2 = -1$, so that the representations in equations (2.2) and (2.3) naturally prompt the notion to view these numbers as a doubling of the complex numbers. The real unit ℓ and the *hyperbolic unit* $k = IJ$ square to 1, $\ell^2 = k^2 = 1$. The multiplication of these units is commutative with further products in the Cayley multiplication table being $\ell I = I, \ell J = J, \ell k = k, Ik = -J, Jk = -I$. The *idempotent representation* (2.5) is an orthogonal decomposition obtained by using the orthogonal idempotents

$$e_1 := \frac{1+k}{2} \quad \text{and} \quad e_2 := \frac{1-k}{2}, \quad (2.6)$$

with properties $e_1^2 = e_1, e_2^2 = e_2, e_1e_2 = 0$ and $e_1 + e_2 = 1$. All four representations (2.2)–(2.5) are uniquely related to each other. For instance, given a bicomplex number in the canonical representation (2.4) in the form

$$n_a = a_1\ell + a_2I + a_3J + a_4k, \quad (2.7)$$

the equivalent representations (2.2), (2.4) and (2.5) are obtained with the identifications

$$\begin{aligned} z_1 &= a_1 + Ia_2, & z_2 &= a_3 + Ia_4, \\ w_1 &= a_1 + Ja_3, & w_2 &= a_2 + Ja_4, \\ v_1^a &= (a_1 + a_4)\ell + (a_2 - a_3)I, & v_2^a &= (a_1 - a_4)\ell + (a_2 + a_3)J. \end{aligned} \quad (2.8)$$

Arithmetic operations are most elegantly and efficiently carried out in the idempotent representation (2.5). For the composition of two arbitrary numbers n_a and n_b we have

$$n_a \circ n_b = v_1^a \circ v_1^b e_1 + v_2^a \circ v_2^b e_2 \quad \text{with } \circ \equiv \pm, \cdot, \div. \quad (2.9)$$

The *hyperbolic numbers* (or *split-complex numbers*) $\mathbb{D} = \{a_1\ell + a_4k \mid a_1, a_4 \in \mathbb{R}\}$ are an important special case of \mathbb{B} obtained in the absence of the imaginary units I and J , or when taking $a_2 = a_3 = 0$.

The same arithmetic rules as in (2.9) then apply to *bicomplex functions*. In what follows we are most interested in functions depending on two real variables x and t of the form $f(x, t) = \ell p(x, t) + Iq(x, t) + Jr(x, t) + ks(x, t) \in \mathbb{B}$ involving four real fields $p(x, t), q(x, t), r(x, t), s(x, t) \in \mathbb{R}$. Having kept the functional variables real, we also keep our differential real, so that we can differentiate $f(x, t)$ componentwise as $\partial_x f(x, t) = \ell \partial_x p(x, t) + I \partial_x q(x, t) + J \partial_x r(x, t) + k \partial_x s(x, t)$ and similarly for $\partial_t f(x, t)$. For further properties of bicomplex numbers and functions, such as for instance computing norms, see for instance [17, 26, 29, 30].

2.2. \mathcal{PT} -symmetric bicomplex functions and conserved quantities

As there are two different imaginary units, there are three different types of conjugations for bicomplex numbers, corresponding to conjugating only I , only J or conjugating both I and J simultaneously.

This is reflected in different symmetries that leave the Cayley multiplication table invariant. As a consequence we also have three different types of bicomplex \mathcal{PT} -symmetries, acting as

$$\mathcal{PT}_{ij} : \ell \rightarrow \ell, \iota \rightarrow -\iota, j \rightarrow -j, k \rightarrow k, x \rightarrow -x, t \rightarrow -t, \tag{2.10}$$

$$\mathcal{PT}_{ik} : \ell \rightarrow \ell, \iota \rightarrow -\iota, j \rightarrow j, k \rightarrow -k, x \rightarrow -x, t \rightarrow -t, \tag{2.11}$$

$$\mathcal{PT}_{jk} : \ell \rightarrow \ell, \iota \rightarrow \iota, j \rightarrow -j, k \rightarrow -k, x \rightarrow -x, t \rightarrow -t, \tag{2.12}$$

see also [3]. When decomposing the bicomplex energy eigenvalue of a bicomplex Hamiltonian H in the time-independent Schrödinger equation, $H\psi = E\psi$, as $E = E_1\ell + E_2\iota + E_3j + E_4k$, Bagchi and Banerjee argued in [3] that a \mathcal{PT}_{ik} -symmetry ensures that $E_2 = E_4 = 0$, a \mathcal{PT}_{jk} -symmetry forces $E_3 = E_4 = 0$ and a \mathcal{PT}_{ij} -symmetry sets $E_2 = E_3 = 0$. In [12–14] we argued that for complex soliton solutions the \mathcal{PT} -symmetries together with the integrability of the model guarantees the reality of all physical conserved quantities. One of the main concerns in this section is to investigate the roles played by the symmetries (2.10)–(2.12) for the bicomplex soliton solutions and to clarify whether the implications are similar as observed in the quantum case.

Decomposing a density function for any conserved quantity as

$$\rho(x, t) = \ell\rho_1(x, t) + \iota\rho_2(x, t) + j\rho_3(x, t) + k\rho_4(x, t) \in \mathbb{B}, \tag{2.13}$$

and demanding it to be \mathcal{PT} -invariant, it is easily verified that a \mathcal{PT}_{ik} -symmetry implies that ρ_1, ρ_3 and ρ_2, ρ_4 are even and odd functions of x , respectively. A \mathcal{PT}_{jk} -symmetry forces ρ_1, ρ_2 and ρ_3, ρ_4 to even and odd in x , respectively and a \mathcal{PT}_{ij} -symmetry makes ρ_1, ρ_4 and ρ_2, ρ_3 even and odd in x , respectively. The corresponding conserved quantities must therefore be of the form

$$Q = \int_{-\infty}^{\infty} \rho(x, t) dx = \begin{cases} Q_1\ell + Q_3j & \text{for } \mathcal{PT}_{ik}\text{-symmetric } \rho \\ Q_1\ell + Q_2\iota & \text{for } \mathcal{PT}_{jk}\text{-symmetric } \rho \\ Q_1\ell + Q_4k & \text{for } \mathcal{PT}_{ij}\text{-symmetric } \rho \end{cases}, \tag{2.14}$$

where we denote $Q_i := \int_{-\infty}^{\infty} \rho_i(x, t) dx$ with $i = 1, 2, 3, 4$. Thus we expect the same property that forces certain quantum mechanical energies to vanish to hold similarly for all classical conserved quantities. We only regard Q_1 and Q_4 as physical, so that only a \mathcal{PT}_{ij} -symmetric system is guaranteed to be physical.

2.3. The bicomplex Korteweg-de Vries equation

Using the multiplication law (2.9) for bicomplex functions, the KdV equation for a bicomplex field in the canonical form

$$u(x, t) = \ell p(x, t) + \iota q(x, t) + jr(x, t) + ks(x, t) \in \mathbb{B}, \tag{2.15}$$

can either be viewed as a set of coupled equations for the four real fields $p(x, t), q(x, t), r(x, t), s(x, t) \in \mathbb{R}$

$$u_t + 6uu_x + u_{xxx} = 0 \Leftrightarrow \begin{cases} p_t + 6pp_x - 6qq_x - 6rr_x + 6ss_x + p_{xxx} = 0 \\ q_t + 6qp_x + 6pq_x - 6sr_x - 6rs_x + q_{xxx} = 0 \\ r_t + 6rp_x + 6pr_x - 6qs_x - 6sq_x + r_{xxx} = 0 \\ s_t + 6sp_x + 6ps_x + 6qr_x + 6rq_x + s_{xxx} = 0 \end{cases}, \tag{2.16}$$

or when using the representation (2.5) as a couple of complex KdV equations

$$v_t + 6vv_x + v_{xxx} = 0, \quad \text{and} \quad w_t + 6ww_x + w_{xxx} = 0, \quad (2.17)$$

related to the canonical representation as

$$v(x, t) = [p(x, t) + s(x, t)] + i[q(x, t) - r(x, t)] \in \mathbb{C}(i), \quad (2.18)$$

$$w(x, t) = [p(x, t) - s(x, t)] + j[q(x, t) + r(x, t)] \in \mathbb{C}(j). \quad (2.19)$$

We recall that we keep here our space and time variables, x and t , to be both real so that also the corresponding derivatives ∂_x and ∂_t are not bicomplexified.

When acting on the component functions the \mathcal{PT} -symmetries (2.10)–(2.12) are implemented in (2.16) as

$$\mathcal{PT}_{ij} : x \rightarrow -x, t \rightarrow -t, p \rightarrow p, q \rightarrow -q, r \rightarrow -r, s \rightarrow s, u \rightarrow u, \quad (2.20)$$

$$\mathcal{PT}_{ik} : x \rightarrow -x, t \rightarrow -t, p \rightarrow p, q \rightarrow -q, r \rightarrow r, s \rightarrow -s, u \rightarrow u, \quad (2.21)$$

$$\mathcal{PT}_{jk} : x \rightarrow -x, t \rightarrow -t, p \rightarrow p, q \rightarrow q, r \rightarrow -r, s \rightarrow -s, u \rightarrow u, \quad (2.22)$$

ensuring that the KdV-equation remains invariant for all of the transformations. Notice that the representation in (2.17) remains only invariant under \mathcal{PT}_{ij} , but does not respect the symmetries \mathcal{PT}_{ik} and \mathcal{PT}_{jk} .

We observe that (2.16) allows for a scaling of space by the hyperbolic unit k as $x \rightarrow kx$, leading to a new type of KdV-equation with $u \rightarrow h$

$$kh_t + 6hh_x + h_{xxx} = 0 \quad \Leftrightarrow \quad \begin{cases} s_t + 6pp_x - 6qq_x - 6rr_x + 6ss_x + p_{xxx} = 0 \\ r_t - 6qp_x - 6pq_x + 6sr_x + 6rs_x - q_{xxx} = 0 \\ q_t - 6rp_x - 6pr_x + 6qs_x + 6sq_x - r_{xxx} = 0 \\ p_t + 6sp_x + 6ps_x + 6qr_x + 6rq_x + s_{xxx} = 0 \end{cases}, \quad (2.23)$$

that also respects the \mathcal{PT}_{ij} -symmetry. The interesting consequence of this modification is that traveling wave solutions $u(\xi)$ of (2.16) depending on real combination of x and t as $\xi = x + ct \in \mathbb{R}$, with c denoting the speed, become solutions $h(\zeta)$ dependent on the hyperbolic number $\zeta = kx + ct \in \mathbb{D}$ instead. Interestingly a hyperbolic rotation of this number ζ , defined as $\zeta' = \zeta e^{-\phi k} = kx' + ct'$ with $\phi = \arctan(v/c)$, constitutes a Lorentz transformation with $t' = \gamma(t - v/c^2 x)$, $x' = \gamma(t - vx)$ and $\gamma = 1/\sqrt{1 - v^2/c^2}$, see e.g. [34, 36].

Next we consider various solutions to these different versions of the bicomplex KdV-equation, discuss how they may be constructed and their key properties.

2.3.1. One-soliton solutions with broken \mathcal{PT} -symmetry

We start from the well known bright one-soliton solution of the real KdV equation (2.16)

$$u_{\mu, \alpha}(x, t) = \frac{\alpha^2}{2} \operatorname{sech}^2 \left[\frac{1}{2}(\alpha x - \alpha^3 t + \mu) \right], \quad (2.24)$$

when $\alpha, \mu \in \mathbb{R}$. Since our differentials have not been bicomplexified we may take μ to be a bicomplex number $\mu = \rho\ell + \theta i + \phi j + \chi k \in \mathbb{B}$ with $\rho, \theta, \phi, \chi \in \mathbb{R}$, so that (2.24) becomes a solution of the bicomplex equation (2.16). We may of course also take α to be bicomplex, but as we commented in [13] already on the complex case, this would imply losing the \mathcal{PT} -symmetry and would also

lead to non real, hence unphysical, conserved quantities. Expanding the hyperbolic function, we can separate the bicomplex function $u_{\mu,\alpha}(x, t)$ after some lengthy computation into its different canonical components

$$u_{\rho,\theta,\phi,\chi;\alpha} = \frac{\ell}{2} [p_{\rho+\chi,\theta-\phi;\alpha} + p_{\rho-\chi,\theta+\phi;\alpha}] + \frac{l}{2} [q_{\rho+\chi,\theta-\phi;\alpha} + q_{\rho-\chi,\theta+\phi;\alpha}] \quad (2.25)$$

$$+ \frac{j}{2} [q_{\rho-\chi,\theta+\phi;\alpha} - q_{\rho+\chi,\theta-\phi;\alpha}] + \frac{k}{2} [p_{\rho+\chi,\theta-\phi;\alpha} - p_{\rho-\chi,\theta+\phi;\alpha}],$$

when using the two functions

$$p_{a,b;\alpha}(x, t) = \frac{\alpha^2 + \alpha^2 \cos b \cosh(\alpha x - \alpha^3 t + a)}{[\cos b + \cosh(\alpha x - \alpha^3 t + a)]^2}, \quad (2.26)$$

$$q_{a,b;\alpha}(x, t) = \frac{\alpha^2 \sin b \sinh(\alpha x - \alpha^3 t + a)}{[\cos b + \cosh(\alpha x - \alpha^3 t + a)]^2}. \quad (2.27)$$

Noting that the complex solution $u_{i\theta,\alpha}(x, t)$ studied in [13], can be expressed as $u_{i\theta,\alpha}(x, t) = p_{a,\theta;\alpha}(x - a/\alpha, t) + iq_{a,\theta;\alpha}(x - a/\alpha, t)$, we can also expand the bicomplex solution (2.25) in terms of the complex solution as

$$u_{\rho,\theta,\phi,\chi;\alpha} = \frac{\ell}{2} \left[u_{i(\phi-\theta),\alpha} \left(x + \frac{\rho+\chi}{\alpha}, t \right) + u_{-i(\phi+\theta),\alpha} \left(x + \frac{\rho-\chi}{\alpha}, t \right) \right] \quad (2.28)$$

$$+ \frac{k}{2} \left[u_{-i(\phi-\theta),\alpha} \left(x + \frac{\rho+\chi}{\alpha}, t \right) - u_{i(\phi+\theta),\alpha} \left(x + \frac{\rho-\chi}{\alpha}, t \right) \right].$$

In figure 2.3.1 we depict the canonical components of this solution at different times. We observe in all of them that the one-soliton solution is split into two separate one-soliton-like components moving parallel to each other with the same speed. The real p -component can be viewed as the sum of two bright solitons and the hyperbolic s -component is the sum of a bright and a dark soliton. This effect is the results of the decomposition of each of the components into a sum of the functions $p_{a,b;\alpha}$ or $q_{a,b;\alpha}$, as defined in (2.26), at different values of a , b , but the same value of α . Since a and b control the amplitude and distance, whereas α regulates the speed, the constituents travel at the same speed. We recall that this type of behaviour of degenerate solitons can neither be created from a real nor a complex two-soliton solution [11, 15]. So this is a novel type of phenomenon for solitons previously not observed.

In general, the solution (2.24) is not \mathcal{PT} -symmetric with regard to any of the possibilities defined above. It becomes \mathcal{PT}_{ij} -symmetric when $\rho = \chi = 0$, \mathcal{PT}_{ik} -symmetric when $\rho = \chi = \phi = 0$ and \mathcal{PT}_{jk} -symmetric when $\rho = \chi = \theta = 0$.

A solution to the new KdV equation (2.23) is constructed as

$$h_{\mu,\alpha}(x, t) = \frac{\alpha^2}{2} \operatorname{sech}^2 \left[\frac{1}{2} (\alpha x k - \alpha^3 t + \mu) \right], \quad (2.29)$$

which in component form reads

$$h_{\rho,\theta,\phi,\chi;\alpha} = \frac{\ell}{2} [\bar{p}_{\chi-\rho,\theta+\phi;\alpha} + p_{\chi+\rho,\theta-\phi;\alpha}] + \frac{l}{2} [\bar{q}_{\chi-\rho,\theta+\phi;\alpha} - q_{\chi+\rho,\theta-\phi;\alpha}] \quad (2.30)$$

$$+ \frac{k}{2} [\bar{q}_{\chi-\rho,\theta+\phi;\alpha} + q_{\chi+\rho,\theta-\phi;\alpha}] + \frac{k}{2} [\bar{p}_{\chi+\rho,\theta-\phi;\alpha} - p_{\chi-\rho,\theta+\phi;\alpha}],$$

where we introduced the notation $\bar{p}_{a,b;\alpha}(x, t) = p_{a,b;\alpha}(x, -t)$ and $\bar{q}_{a,b;\alpha}(x, t) = q_{a,b;\alpha}(x, -t)$.




Fig. 1. Canonical component functions p , q , r and s (clockwise starting in the top left corner) of the decomposed one-soliton solution $u_{\rho,\theta,\phi,\chi;\alpha}$ to the bicomplex KdV equation (2.16) with broken \mathcal{PT} -symmetry at different times for $\alpha = 0.5$, $\rho = 1.3$, $\theta = 0.4$, $\phi = 2.0$ and $\chi = 1.3$.

In figure 2.3.1 we depict the canonical component functions of this solution. We observe that the one-soliton solution is split into two one-soliton-like structures that scatter head-on with each other. The real p -component consists of a head-on scattering of two bright solitons and hyperbolic the s -component is a head-on collision of a bright and a dark soliton. Given that $u_{\rho,\theta,\phi,\chi;\alpha}$ and $h_{\rho,\theta,\phi,\chi;\alpha}(x,t)$ differ in the way that one of its constituent functions is time-reversed this is to be expected.

2.3.2. \mathcal{PT}_{ij} -symmetric one-soliton solution

An interesting solution can be constructed when we start with a complex \mathcal{PT}_{ik} and a complex \mathcal{PT}_{jk} symmetric solution to assemble the linear decomposition of an overall \mathcal{PT}_{ij} -symmetric solution with different velocities. Taking in the decomposition (2.17) $v(x,t) = u_{i\theta,\alpha}(x,t)$ and $w(x,t) = u_{j\phi,\beta}(x,t)$, we can build the bicomplex KdV-solution in the idempotent representation

$$\hat{u}_{\theta,\phi;\alpha,\beta}(x,t) = u_{i\theta,\alpha}(x,t)e_1 + u_{j\phi,\beta}(x,t)e_2. \quad (2.31)$$

The expanded version in the canonical representation becomes in this case

$$\hat{u}_{\theta,\phi;\alpha,\beta} = \frac{\ell}{2} [p_{0,\theta;\alpha} + p_{0,\phi;\beta}] + \frac{l}{2} [q_{0,\theta;\alpha} + q_{0,\phi;\beta}] + \frac{j}{2} [q_{0,\phi;\beta} - q_{0,\theta;\alpha}] + \frac{k}{2} [p_{0,\theta;\alpha} - p_{0,\phi;\beta}], \quad (2.32)$$




Fig. 2. Head-on collision of a bright soliton with a dark soliton in the canonical components p, q, r, s (clockwise starting in the top left corner) for the one-soliton solution $h_{\rho, \theta, \phi, \chi; \alpha}$ to the bicomplex KdV equation (2.16) with broken \mathcal{PT} -symmetry for $\alpha = 0.5, \rho = 1.3, \theta = 0.1, \phi = 2.0$ and $\chi = 1.3$. Time is running vertically, space horizontally and contours of the amplitudes are colour-coded indicated as in the legends.

which is evidently \mathcal{PT}_{IJ} -symmetric. Hence this solution contain any multicomplex shifts, but in each component two solitonic contributions with different amplitude and speed parameter. As we can see in figure 2.3.2, in the real p -component a faster bright soliton is overtaking a slower bright solitons and in hyperbolic s -component a faster bright soliton is overtaking and a slower dark soliton. Unlike as in the real or complex case, one can carry out the limit $\beta \rightarrow \alpha$ to the degenerate case without complication since have the identity $\hat{u}_{\theta-\phi, \theta+\phi; \alpha, \alpha} = u_{0, \theta, \phi, 0; \alpha}$. Similarly as in the previous section we may also construct a further solution from a hyperbolic shift $x \rightarrow kx$, which we do not present here.

2.3.3. Multi-soliton solutions

The most compact way to express the N -soliton solution for the real KdV equation in the form (2.16) is

$$u_{\mu_1, \mu_2, \dots, \mu_n; \alpha_1, \alpha_2, \dots, \alpha_n}^{(n)}(x, t) = 2 \left[\ln W_n(\psi_{\mu_1, \alpha_1}, \psi_{\mu_2, \alpha_2}, \dots, \psi_{\mu_n, \alpha_n}) \right]_{xx}, \quad (2.33)$$




Fig. 3. A fast bright soliton overtaking a slower bright soliton in the canonical component functions p , q , r and s (clockwise starting in the top left corner) for the one-soliton solution $\hat{u}_{\theta,\phi;\alpha,\beta}$ to the bicomplex KdV equation (2.16) with \mathcal{PT}_{ij} -symmetry for $\alpha = 2.1$, $\beta = 1.1$, $\theta = 0.6$ and $\phi = 1.75$.

where $W_n[\psi_1, \psi_2, \dots, \psi_n] := \det \omega$ denotes the Wronskian with $\omega_{jk} = \partial^{j-1} \psi_k / \partial x^{j-1}$ for $j, k = 1, \dots, n$, e.g. $W_1[\psi_0] = \psi_0$, $W_2[\psi_0, \psi_1] = \psi_0(\psi_1)_x - \psi_1(\psi_0)_x$, etc and the functions ψ_i are solutions to the time-independent Schrödinger equation for the free theory. Taking for instance $\psi_{\mu,\alpha}(x, t) = \cosh[(\alpha x - \alpha^3 t + \mu)/2]$ for $n = 1$ leads to the one-soliton solution (2.24).

We could now take the shifts $\mu_1, \mu_2, \dots, \mu_n \in \mathbb{B}$ and expand (2.33) into its canonical components to obtain the N -soliton solution for the bicomplex equation. Alternatively we may also construct N -solitons in the idempotent basis in analogy to (2.32). We demonstrate here the latter approach for the two-soliton. From (2.33) we observe that the second derivative will not alter the linear bicomplex decomposition and it is therefore useful to introduce the quantity $w(x, t)$ as $u = w_x$. Thus a complex one-soliton solution can be obtained from

$$w_{a,b;\alpha}(x, t) = w_{a,b;\alpha}^r(x, t) + \iota w_{a,b;\alpha}^i(x, t) \quad (2.34)$$

with

$$w_{a,b;\alpha}^r(x,t) = \frac{\alpha \sinh(\alpha x - \alpha^3 t + a)}{\cos b + \cosh(\alpha x - \alpha^3 t + a)}, \quad w_{a,b;\alpha}^i(x,t) = \frac{\alpha \sin b}{\cos b + \cosh(\alpha x - \alpha^3 t + a)}. \quad (2.35)$$

Noting that $p_{a,b;\alpha} = (w_{a,b;\alpha}^r)_x$, $q_{a,b;\alpha} = (w_{a,b;\alpha}^i)_x$ we obtain a complex soliton as $u_{a,b;\alpha} = (w_{a,b;\alpha})_x$. Recalling now the expression

$$w_{a,b,c,d;\alpha,\beta} = \frac{\alpha^2 - \beta^2}{w_{a,b;\alpha} - w_{c,d;\beta}}, \quad (2.36)$$

from the Bäcklund transformation of the complex two-soliton [13], we can express this in terms of the functions in (2.35)

$$w_{a,b,c,d;\alpha,\beta} = \frac{(\alpha^2 - \beta^2) \left[(w_{a,b;\alpha}^r - w_{c,d;\beta}^r) - \iota (w_{a,b;\alpha}^i - w_{c,d;\beta}^i) \right]}{(w_{a,b;\alpha}^r - w_{c,d;\beta}^r)^2 + (w_{a,b;\alpha}^i - w_{c,d;\beta}^i)^2} = w_{a,b,c,d;\alpha,\beta}^r + \iota w_{a,b,c,d;\alpha,\beta}^i. \quad (2.37)$$

Using (2.37) to define the two complex quantities $w_{\theta_1, \theta_2, \theta_3, \theta_4; \alpha_1, \alpha_2} = w_2^r + \iota w_2^i \in \mathbb{C}(\iota)$ and $w_{\phi_1, \phi_2, \phi_3, \phi_4; \beta_1, \beta_2} = \tilde{w}_2^r + J \tilde{w}_2^i \in \mathbb{C}(J)$ we introduce the bicomplex function

$$w_2 = (w_2^r + \iota w_2^i) e_1 + (\tilde{w}_2^r + J \tilde{w}_2^i) e_2 \quad (2.38)$$

$$= \frac{\ell}{2} (w_2^r + \tilde{w}_2^r) + \frac{\iota}{2} (w_2^i + \tilde{w}_2^i) + \frac{J}{2} (\tilde{w}_2^i - w_2^i) + \frac{k}{2} (w_2^r - \tilde{w}_2^r). \quad (2.39)$$

Then by construction $u_{\theta_1, \theta_2, \theta_3, \theta_4, \phi_1, \phi_2, \phi_3, \phi_4; \alpha_1, \alpha_2, \beta_1, \beta_2} = (w_2)_x$ is a bicomplex two-soliton solution with four speed parameters. In a similar fashion we can proceed to construct N -soliton for $N > 2$.

2.3.4. Real and hyperbolic conserved quantities

Next we compute the first conserved quantities the mass m , the momentum p and the energy E , see e.g. [12, 13]

$$m(u) = \int_{-\infty}^{\infty} u dx = m_1 \ell + m_2 \iota + m_3 J + m_4 k, \quad (2.40)$$

$$p(u) = \int_{-\infty}^{\infty} u^2 dx = p_1 \ell + p_2 \iota + p_3 J + p_4 k, \quad (2.41)$$

$$E(u) = \int_{-\infty}^{\infty} \left(\frac{1}{2} u_x^2 - u^3 \right) dx = E_1 \ell + E_2 \iota + E_3 J + E_4 k \quad (2.42)$$

Decomposing the relevant densities into the canonical basis, u as in (2.15), u^2 as

$$u^2 = (p^2 - q^2 - r^2 + s^2) \ell + 2(pq - rs) \iota + 2(pr - qs) J + 2(qr + ps) k \quad (2.43)$$

and the Hamiltonian density $\mathcal{H}(u, u_x) = u_x^2/2 - u^3$ as

$$\begin{aligned} \mathcal{H} = & \left[3p \left(q^2 + r^2 - s^2 \right) + \frac{p_x^2 - q_x^2 - r_x^2 + s_x^2}{2} - 6qrs - p^3 \right] \ell \\ & + \left[q^3 - 3p^2q + p_xq_x + 6prs + 3q \left(r^2 - s^2 \right) - r_x s_x \right] \iota \\ & + \left[r^3 + 6pqs + 3r \left(q^2 - s^2 - p^2 \right) + p_x r_x - q_x s_x \right] j \\ & + \left[3s \left(r^2 - p^2 + q^2 \right) - 6pqr + p_x s_x + q_x r_x - s^3 \right] k, \end{aligned} \quad (2.44)$$

we integrate componentwise. For the solutions $u_{\rho, \theta, \phi, \chi; \alpha}$ and $h_{\rho, \theta, \phi, \chi; \alpha}$ with broken \mathcal{PT} -symmetry we obtain the real conserved quantities

$$m(u_{\rho, \theta, \phi, \chi; \alpha}) = m(h_{\rho, \theta, \phi, \chi; \alpha}) = 2\alpha\ell, \quad (2.45)$$

$$p(u_{\rho, \theta, \phi, \chi; \alpha}) = p(h_{\rho, \theta, \phi, \chi; \alpha}) = \frac{2}{3}\alpha^3\ell, \quad (2.46)$$

$$E(u_{\rho, \theta, \phi, \chi; \alpha}) = E(h_{\rho, \theta, \phi, \chi; \alpha}) = -\frac{1}{5}\alpha^5\ell. \quad (2.47)$$

These values are the same as those found in [13] for the complex solitons. Given that the \mathcal{PT} -symmetries are all broken this is surprising at first sight. However, considering the representation (2.28) this is easily understood when using the result of [13]. Then $m(u_{\rho, \theta, \phi, \chi; \alpha})$ is simply $\ell/2(2\alpha + 2\alpha) + j/2(2\alpha - 2\alpha) = 2\alpha\ell$. We can argue similarly for the other conserved quantities.

For the \mathcal{PT}_{ij} -symmetric solution $\hat{u}_{\theta, \phi; \alpha, \beta}$ we obtain the following hyperbolic values for the conserved quantities

$$m(\hat{u}_{\theta, \phi; \alpha, \beta}) = (\alpha + \beta)\ell + (\alpha - \beta)k, \quad (2.48)$$

$$p(\hat{u}_{\theta, \phi; \alpha, \beta}) = \frac{1}{3}(\alpha^3 + \beta^3)\ell + \frac{1}{3}(\alpha^3 - \beta^3)k \quad (2.49)$$

$$E(\hat{u}_{\theta, \phi; \alpha, \beta}) = -\left(\frac{\alpha^5}{10} + \frac{\beta^5}{10}\right)\ell + \left(\frac{\beta^5}{10} - \frac{\alpha^5}{10}\right)k. \quad (2.50)$$

The values become real and coincide with the expressions (2.45)–(2.47) when we sum up the contributions from the real and hyperbolic component or in the degenerate case when we take the limit $\beta \rightarrow \alpha$.

2.4. The bicomplex Alice and Bob KdV equation

Various nonlocal versions of nonlinear wave equations that have been overlooked previously have attracted considerable attention recently. In reference to standard scenarios in quantum cryptography some of them are also often referred to as Alice and Bob systems. These variants of the nonlinear Schrödinger or Hirota equation [1, 10, 27, 33] arise from an alternative choice in the compatibility condition of the two AKNS-equations. For the KdV equation (2.16) they can be constructed [23–25] by choosing $u(x, t) = 1/2[a(x, t) + b(x, t)]$, with the constraint $\mathcal{PT}a(x, t) = a(-x, -t) = b(x, t)$, thus converting it into an equation that can be decomposed into two equations, the Alice and Bob KdV

(ABKdV) equation

$$a_t + 3/4(a + b)(3a_x + b_x) + a_{xxx} = 0, \quad (2.51)$$

$$b_t + 3/4(a + b)(a_x + 3b_x) + b_{xxx} = 0. \quad (2.52)$$

In a similar way as the two AKNS-equations can be made compatible by a suitable transformation map, these two equations are converted into each other by a \mathcal{PT} -transformation, i.e. $\mathcal{PT}(2.51) \equiv (2.52)$. Evidently the decomposition is not unique and one may also add and subtract a constrained function of a and b or consider different types of maps to relate the equation.

The bicomplex version of the Alice and Bob system (2.51), (2.52) is obtained by taking $a, b \in \mathbb{B}$. In the canonical basis we use the conventions $u(x, t) = \ell p(x, t) + iq(x, t) + jr(x, t) + ks(x, t)$, $a(x, t) = \ell \hat{p}(x, t) + i\hat{q}(x, t) + j\hat{r}(x, t) + k\hat{s}(x, t)$, $b(x, t) = \ell \check{p}(x, t) + i\check{q}(x, t) + j\check{r}(x, t) + k\check{s}(x, t)$, so that the ABKdV equations (2.51) and (2.52) decompose into eight coupled equations

$$\hat{p}_t = -\hat{p}_{xxx} - \frac{3}{2} [p(\check{p}_x + 3\hat{p}_x) - q(\check{q}_x + 3\hat{q}_x) - r(\check{r}_x + 3\hat{r}_x) + s(\check{s}_x + 3\hat{s}_x)], \quad (2.53)$$

$$\hat{q}_t = -\hat{q}_{xxx} + \frac{3}{2} [p(\check{q}_x + 3\hat{q}_x) + q(\check{p}_x + 3\hat{p}_x) - r(\check{s}_x + 3\hat{s}_x) - s(\check{r}_x + 3\hat{r}_x)], \quad (2.54)$$

$$\hat{r}_t = -\hat{r}_{xxx} + \frac{3}{2} [p(\check{r}_x + 3\hat{r}_x) - q(\check{s}_x + 3\hat{s}_x) + r(\check{p}_x + 3\hat{p}_x) - s(\check{q}_x + 3\hat{q}_x)], \quad (2.55)$$

$$\hat{s}_t = -\hat{s}_{xxx} + \frac{3}{2} [p(\check{s}_x + 3\hat{s}_x) + q(\check{r}_x + 3\hat{r}_x) + r(\check{q}_x + 3\hat{q}_x) + s(\check{p}_x + 3\hat{p}_x)], \quad (2.56)$$

and

$$\check{p}_t = -\check{p}_{xxx} + \frac{3}{2} [p(3\check{p}_x + \hat{p}_x) - q(3\check{q}_x + \hat{q}_x) - r(3\check{r}_x + \hat{r}_x) + s(3\check{s}_x + \hat{s}_x)], \quad (2.57)$$

$$\check{q}_t = -\check{q}_{xxx} + \frac{3}{2} [p(3\check{q}_x + \hat{q}_x) + q(3\check{p}_x + \hat{p}_x) - r(3\check{s}_x + \hat{s}_x) - s(3\check{r}_x + \hat{r}_x)], \quad (2.58)$$

$$\check{r}_t = -\check{r}_{xxx} + \frac{3}{2} [p(3\check{r}_x + \hat{r}_x) - q(3\check{s}_x + \hat{s}_x) + r(3\check{p}_x + \hat{p}_x) - s(3\check{q}_x + \hat{q}_x)], \quad (2.59)$$

$$\check{s}_t = -\check{s}_{xxx} + \frac{3}{2} [p(3\check{s}_x + \hat{s}_x) + q(3\check{r}_x + \hat{r}_x) + r(3\check{q}_x + \hat{q}_x) + s(3\check{p}_x + \hat{p}_x)]. \quad (2.60)$$

A real solution to the ABKdV equations (2.51) and (2.52) that sums up to the standard one-soliton solution (2.24) is found as

$$a_{\mu, \nu; \alpha}(x, t) = u_{\mu, \alpha}(x, t) + \nu \tanh \left[\frac{1}{2}(\alpha x - \alpha^3 t + \mu) \right], \quad (2.61)$$

$$b_{\mu, \nu; \alpha}(x, t) = u_{\mu, \alpha}(x, t) - \nu \tanh \left[\frac{1}{2}(\alpha x - \alpha^3 t + \mu) \right], \quad (2.62)$$

with arbitrary constants $\nu, \mu \in \mathbb{R}$. Proceeding now as for the local variant by taking $\mu = \rho\ell + \theta t + \phi J + \chi k \in \mathbb{B}$, we decompose $a_{\mu, \nu; \alpha}$ and $b_{\mu, \nu; \alpha}$ into their canonical components and obtain after some lengthy computation the corresponding solution to the bicomplex version of the ABKdV equations

(2.53)–(2.60) as

$$a_{\rho,\theta,\phi,\chi;\alpha} = u_{\rho,\theta,\phi,\chi;\alpha} + \frac{\ell}{2}vF_{w+\rho,\theta,\phi,\chi} + \frac{l}{2}vG_{w+\rho,\theta,\phi,\chi} + \frac{J}{2}vG_{w+\rho,\theta,\phi,\chi} + \frac{k}{2}vF_{\chi,\theta,\phi,w+\rho} \quad (2.63)$$

with $w_\alpha = \alpha x - \alpha^3 t$ and the newly defined functions

$$F_{x_1,x_2,x_3,x_4} = \frac{\sinh x_1 \sec x_2 \sec x_3 \operatorname{sech} x_4 + \tanh x_1 - \tan x_2 \tan x_3 \tanh x_4}{1 - \tanh x_1 \tan x_2 \tan x_3 \tanh x_4 + \frac{\cosh(2x_1)+\cos(2x_2)+\cos(2x_3)+\cosh(2x_4)}{4 \cosh x_1 \cos x_2 \cos x_3 \cosh x_4}}, \quad (2.64)$$

$$G_{x_1,x_2,x_3,x_4} = \frac{\operatorname{sech} x_1 \sec x_2 \sin x_3 \operatorname{sech} x_4 + \tan x_3 + \tanh x_1 \tan x_2 \tanh x_4}{1 - \tanh x_1 \tan x_2 \tan x_3 \tanh x_4 + \frac{\cosh(2x_1)+\cos(2x_2)+\cos(2x_3)+\cosh(2x_4)}{4 \cosh x_1 \cos x_2 \cos x_3 \cosh x_4}}. \quad (2.65)$$

The functions $b_{\rho,\theta,\phi,\chi;\alpha}$, or equivalently the individual components $\check{p}, \check{q}, \check{r}, \check{s}$, are obtained by a \mathcal{PT} -transformation.

We may also proceed as in subsection 2.3.2 and construct a solution in the idempotent representation. Keeping the parameter v real, a solution based on the idempotent decomposition is

$$\begin{aligned} a_{\theta,\phi,v;\alpha,\beta} &= a_{i\theta,v;\alpha}e_1 + a_{i\phi,v;\beta}e_2 \quad (2.66) \\ &= \hat{u}_{\theta,\phi;\alpha,\beta} + \frac{\ell}{2}v(F_{w_\alpha,\theta,0,0} + F_{w_\beta,\phi,0,0}) + \frac{l}{2}v(G_{w_\alpha,0,\theta,0} + G_{w_\beta,0,\phi,0}) \\ &\quad + \frac{J}{2}v(G_{w_\beta,0,\phi,0} + G_{w_\alpha,0,\theta,0}) + \frac{k}{2}v(F_{w_\alpha,\theta,0,0} - F_{w_\beta,\phi,0,0}). \end{aligned}$$

Once more, the functions $b_{\rho,\theta,\phi,\chi;\alpha}$ or $\check{p}, \check{q}, \check{r}, \check{s}$ are obtained by a \mathcal{PT} -transformation. Comparing (2.66) with $a_{\rho,\theta,\phi,\chi;\alpha}$ in (2.63) we have now two speed parameters at our disposal, similarly as in the local case.

3. Quaternionic solitons

3.1. Quaternionic numbers and functions

The quaternions in the canonical basis are defined as the set of elements

$$\mathbb{H} = \{a_1\ell + a_2i + a_3J + a_4k \mid a_1, a_2, a_3, a_4 \in \mathbb{R}\}. \quad (3.1)$$

The multiplication of the basis $\{\ell, i, J, k\}$ is noncommutative with ℓ denoting the real unit element, $\ell^2 = 1$ and i, J, k its three imaginary units with $i^2 = j^2 = k^2 = -1$. The remaining multiplication rules are $ij = -ji = k$, $jk = -kj = i$ and $kl = -lk = j$. The multiplication table remains invariant under the symmetries \mathcal{PT}_{ij} , \mathcal{PT}_{ik} and \mathcal{PT}_{jk} . Using these rules for the basis, two quaternions in the canonical basis $n_a = a_1\ell + a_2i + a_3J + a_4k \in \mathbb{H}$ and $n_b = b_1\ell + b_2i + b_3J + b_4k \in \mathbb{H}$ are multiplied as

$$\begin{aligned} n_a n_b &= (a_1 b_1 - a_2 b_2 - a_3 b_3 - a_4 b_4)\ell + (a_1 b_2 + a_2 b_1 + a_3 b_4 - a_4 b_3)i \\ &\quad + (a_1 b_3 - a_2 b_4 + a_3 b_1 + a_4 b_2)J + (a_1 b_4 + a_2 b_3 - a_3 b_2 + a_4 b_1)k. \end{aligned} \quad (3.2)$$

There are various representations for quaternions, see e.g. [31], of which the complex form will be especially useful for what follows. With the help of (3.2) one easily verifies that

$$\xi := \frac{1}{\mathcal{N}}(a_2i + a_3J + a_4k) \quad \text{with } \mathcal{N} = \sqrt{a_2^2 + a_3^2 + a_4^2} \quad (3.3)$$

constitutes a new imaginary unit with $\xi^2 = -1$. This means that in this representation we can formally view a quaternion, $n_a \in \mathbb{H}$, as an element in the complex numbers

$$n_a = a_1 \ell + \xi \mathcal{N} \in \mathbb{C}(\xi), \tag{3.4}$$

with real part a_1 and imaginary part \mathcal{N} . Notice that a \mathcal{PT}_{ξ} -symmetry can only be achieved with a \mathcal{PT}_{ijk} -symmetry acting on the unit vectors in the canonical representation. Unlike the bicomplex numbers or the coquaternions, see below, the quaternionic algebra does not contain any idempotents.

3.2. The quaternionic Korteweg-de Vries equation

Applying now the multiplication law (3.2) to quaternionic functions, the KdV equation for a quaternionic field of the form $u(x, t) = \ell p(x, t) + iq(x, t) + jr(x, t) + ks(x, t) \in \mathbb{H}$ can also be viewed as a set of coupled equations for the four real fields $p(x, t), q(x, t), r(x, t), s(x, t) \in \mathbb{R}$

$$u_t + 6uu_x + u_{xxx} = 0 \iff \begin{cases} p_t + 6pp_x - 6qq_x - 6rr_x - 6ss_x + p_{xxx} = 0 \\ q_t + 6qp_x + 6pq_x - 6sr_x + 6rs_x + q_{xxx} = 0 \\ r_t + 6rp_x + 6pr_x - 6qs_x + 6sq_x + r_{xxx} = 0 \\ s_t + 6sp_x + 6ps_x + 6qr_x - 6rq_x + s_{xxx} = 0 \end{cases}. \tag{3.5}$$

Notice that when comparing the bicomplex KdV equation (2.16) and the quaternionic KdV equation (3.5) only the signs of the penultimate terms in all four equations have changed. This means that also (3.5) is invariant under the \mathcal{PT}_{ij} -symmetry. Alternatively, we may consider here the aforementioned symmetry

$$\mathcal{PT}_{ijk} : x \rightarrow -x, t \rightarrow -t, i \rightarrow -i, j \rightarrow -j, k \rightarrow -k, p \rightarrow p, q \rightarrow -q, r \rightarrow -r, s \rightarrow -s, u \rightarrow u, \tag{3.6}$$

which violates all the noncommutative multiplication rules $ij = -ji = k, jk = -kj = i$ and $ki = -ik = j$. Thus in order to implement the symmetry \mathcal{PT}_{ijk} we must set all terms resulting from these multiplications to zero, so that we obtain the additional constraints

$$sr_x = rs_x, \quad qs_x = sq_x, \quad \text{and} \quad qr_x = rq_x. \tag{3.7}$$

When eliminating these terms from (3.5) the remaining set of equations is \mathcal{PT}_{ijk} -symmetric, which appears to be a rather strong imposition. However, the equations without these terms emerge quite naturally when keeping in mind that the product of functions in (3.5) is noncommutative so that one should symmetrize products and replace $6uu_x \rightarrow 3uu_x + 3u_xu$. This process corresponds precisely to imposing the constraints (3.7).

3.3. \mathcal{PT}_{ijk} -symmetric N -soliton solutions

Due to the noncommutative nature of the quaternions it appears difficult at first sight to find solutions to the quaternionic KdV equation. However, using the complex representation (3.4), and imposing the \mathcal{PT}_{ijk} -symmetric, we may resort to our previous analysis on complex solitons. Following [13] and considering the shifted solution (2.24) in the complex space $\mathbb{C}(\xi)$ yields the solution

$$u_{a_1 \ell + \xi \mathcal{N}, \alpha}(x, t) = p_{a_1, \mathcal{N}; \alpha}(x, t) - \xi q_{a_1, \mathcal{N}; \alpha}(x, t) \tag{3.8}$$

$$= p_{a_1, \mathcal{N}; \alpha}(x, t) \ell - \frac{1}{\mathcal{N}} q_{a_1, \mathcal{N}; \alpha}(x, t) (a_2 \ell + a_3 j + a_4 k). \tag{3.9}$$

This solution becomes \mathcal{PT}_{ijk} -symmetric when we carry out a shift in x or t to eliminate the real part of the shift. Reading off the functions $p(x, t), q(x, t), r(x, t), s(x, t)$ from (3.9), it is also obvious

that the constraints (3.7) are indeed satisfied. Thus the real ℓ -component is a one-solitonic structure similar to the real part of a complex soliton and the remaining component consists of the imaginary parts of a complex soliton with overall different amplitudes. It is clear that the conserved quantities constructed from this solution must be real, which follows by using the same argument as for the imaginary part in the complex case [13] separately for each of the i, j, k -components. By considering all functions to be in $\mathbb{C}(\xi)$, it is also clear that multi-soliton solutions can be constructed in analogy to the complex case $\mathbb{C}(t)$ treated in [13] with a subsequent expansion into canonical components.

Since the quaternionic algebra does not contain any idempotents, a construction similar to the one carried out in subsection 2.3.2 does not seem to be possible for quaternions. However, we can use (2.36) for two complex solutions $w_{a,b;\alpha}(x,t) = w_{a,b;\alpha}^r(x,t) + \xi_\alpha w_{a,b;\alpha}^i(x,t)$, $w_{c,d;\beta}(x,t) = w_{c,d;\beta}^r(x,t) + \xi_\beta w_{c,d;\beta}^i(x,t)$, where the imaginary units are defined as in (3.3) with $\xi_a(a_2, a_3, a_4)$ and $\xi_b(b_2, b_3, b_4)$. Expanding that expression in the canonical basis we obtain

$$w_2 = \frac{\alpha^2 - \beta^2}{\omega_1^2 + \omega_2^2 + \omega_3^2 + \omega_4^2} (\ell\omega_1 - i\omega_2 - j\omega_3 - k\omega_4) \quad (3.10)$$

with

$$\omega_1 = w_{a,b;\alpha}^r - w_{c,d;\beta}^r, \quad \omega_n = \frac{a_n w_{a,b;\alpha}^i}{N_a} - \frac{b_n w_{c,d;\beta}^i}{N_b}, \quad n = 2, 3, 4. \quad (3.11)$$

A coquaternionic two-soliton solution to (3.5) is then obtained from (3.10) as $u^{(2)} = (w_2)_x$.

4. Coquaternionic solitons

4.1. Coquaternionic numbers and functions

The coquaternions or often also referred to as split-quaternions in the canonical basis are defined as the set of elements

$$\mathbb{P} = \{a_1 \ell + a_2 i + a_3 j + a_4 k \mid a_1, a_2, a_3, a_4 \in \mathbb{R}\}. \quad (4.1)$$

The multiplication of the basis $\{\ell, i, j, k\}$ is noncommutative with a real unit element ℓ , $\ell^2 = 1$, two hyperbolic unit elements j, k , $j^2 = k^2 = 1$, and one imaginary unit $i^2 = -1$. The remaining multiplication rules are $ij = -ji = k$, $jk = -kj = -i$ and $ki = -ik = j$. The multiplication table remains invariant under the symmetries \mathcal{PT}_{ij} , \mathcal{PT}_{ik} and \mathcal{PT}_{jk} . Using these rules for the basis, two coquaternions in the canonical basis $n_a = a_1 \ell + a_2 i + a_3 j + a_4 k \in \mathbb{P}$ and $n_b = b_1 \ell + b_2 i + b_3 j + b_4 k \in \mathbb{P}$ are multiplied as

$$\begin{aligned} n_a n_b &= (a_1 b_1 - a_2 b_2 + a_3 b_3 + a_4 b_4) \ell + (a_1 b_2 + a_2 b_1 - a_3 b_4 + a_4 b_3) i \\ &+ (a_1 b_3 - a_2 b_4 + a_3 b_1 + a_4 b_2) j + k (a_1 b_4 + a_2 b_3 - a_3 b_2 + a_4 b_1). \end{aligned} \quad (4.2)$$

There are various coquaternionic representations for numbers and functions. Similar as a quaternion one can formally view a coquaternion, $n_1 \in \mathbb{P}$, as an element in the complex numbers

$$n_a = a_1 \ell + \zeta \mathcal{M} \in \mathbb{C}(\zeta) \quad (4.3)$$

with real part a_1 and imaginary part \mathcal{M} . The new imaginary unit, $\zeta^2 = -1$,

$$\zeta := \frac{1}{\mathcal{M}} (a_2 i + a_3 j + a_4 k) \quad \text{with } \mathcal{M} = \sqrt{a_2^2 - a_3^2 - a_4^2} \quad (4.4)$$

is, however, only defined for $a_2^2 \neq a_3^2 + a_4^2$. For definiteness we assume here $|a_2| > \sqrt{a_3^2 + a_4^2}$. Similarly as the \mathcal{PT}_ξ -symmetry also the \mathcal{PT}_ζ -symmetry requires a \mathcal{PT}_{ijk} -symmetry. Unlike the quaternions, the coquaternions possess a number idempotents $e_1 = (1+k)/2$, $e_2 = (1-k)/2$ with $e_1^2 = e_1$, $e_2^2 = e_2$, $e_1e_2 = 0$ or $e_3 = (1+j)/2$, $e_4 = (1-j)/2$ with $e_3^2 = e_3$, $e_4^2 = e_4$, $e_3e_4 = 0$. So for instance, n_a is an element in

$$\mathbb{P} = \{e_1v_1 + e_2v_2 \mid v_1 \in \mathbb{D}(j), v_2 \in \mathbb{D}(j)\}, \quad (4.5)$$

where the hyperbolic numbers in (4.5) are related to the coefficient in the canonical basis as $v_1 = (a_1 + a_4)\ell + (a_2 + a_3)j$ and $v_2 = (a_1 - a_4)\ell + (a_3 - a_2)j$.

4.2. The coquaternionic Korteweg-de Vries equation

Applying now the multiplication law (4.2) to coquaternionic functions, the KdV equation for a quaternionic field of the form $u(x, t) = \ell p(x, t) + iq(x, t) + jr(x, t) + ks(x, t) \in \mathbb{P}$ can also be viewed as a set of coupled equations for the four real fields $p(x, t)$, $q(x, t)$, $r(x, t)$, $s(x, t) \in \mathbb{R}$. The symmetric coquaternionic KdV equation then becomes

$$u_t + 3(uu_x + u_xu) + u_{xxx} = 0 \Leftrightarrow \begin{cases} p_t + 6pp_x - 6qq_x + 6ss_x + 6rr_x + p_{xxx} = 0 \\ q_t + 6qp_x + 6pq_x + q_{xxx} = 0 \\ r_t + 6rp_x + 6pr_x + r_{xxx} = 0 \\ s_t + 6sp_x + 6ps_x + s_{xxx} = 0 \end{cases}. \quad (4.6)$$

Notice that the last three equations of the coupled equation in (4.6) are identical to the symmetric quaternionic KdV equation (3.5) with constraints (3.7).

4.3. \mathcal{PT}_{ijk} -symmetric N -soliton solutions

Using the representation (4.3) we proceed as in subsection 3.3 and consider the shifted solution (2.24) in the complex space $\mathbb{C}(\zeta)$

$$u_{a_1\ell + \zeta M, \alpha}(x, t) = p_{a_1, M; \alpha}(x, t) - \zeta q_{a_1, M; \alpha}(x, t) \quad (4.7)$$

$$= p_{a_1, M; \alpha}(x, t)\ell - \frac{1}{M}q_{a_1, M; \alpha}(x, t)(a_2i + a_3j + a_4k) \quad (4.8)$$

that solves the coquaternionic KdV equation (4.6). The solution in (4.7) is \mathcal{PT}_{ijk} -symmetric. Multi-soliton solutions can be constructed in analogy to the complex case $\mathbb{C}(t)$ treated in [13] by treating all functions in $\mathbb{C}(\zeta)$ as explained in more detail at the end of section 4.

5. Octonionic solitons

We finish our discussion with a comment on the construction of octonionic solitons. Octonions or Cayley numbers are extensions of the quaternions with a doubling of the dimensions. In the canonical basis they can be represented as

$$\mathbb{O} = \{a_0e_0 + a_1e_1 + a_2e_2 + a_3e_3 + a_4e_4 + a_5e_5 + a_6e_6 + a_7e_7 \mid a_i \in \mathbb{R}\}. \quad (5.1)$$

The multiplication of the units is defined by noting that each of the seven quadruplets (e_0, e_1, e_2, e_3) , (e_0, e_1, e_4, e_5) , (e_0, e_1, e_7, e_6) , (e_0, e_2, e_4, e_6) , (e_0, e_2, e_5, e_7) , (e_0, e_3, e_4, e_7) and (e_0, e_3, e_6, e_5) , constitutes a

canonical basis for the quaternions in one-to-one correspondence with (ℓ, i, j, k) . Hence the octonions have one real unit, 7 imaginary units and the multiplication of two octonions is noncommutative. Similarly as for quaternions and coquaternions we can view an octonion $n_a \in \mathbb{O}$ as a complex number

$$n_a = a_1 \ell + o \mathcal{O} \in \mathbb{C}(o) \quad (5.2)$$

with real part a_1 , imaginary part \mathcal{O} and newly defined imaginary unit, $o^2 = -1$,

$$o := \frac{1}{\mathcal{O}} \sum_{i=1}^7 a_i e_i \quad \text{where } \mathcal{O} = \sqrt{\sum_{i=1}^7 a_i e_i}. \quad (5.3)$$

In order to obtain a \mathcal{PT}_o -symmetry we require a $\mathcal{PT}_{e_1 e_2 e_3 e_4 e_5 e_6 e_7}$ -symmetry in the canonical basis.

5.1. The octonionic Korteweg-de Vries equation

Taking now an octonionic field to be of the form $u(x, t) = p(x, t)e_0 + q(x, t)e_1 + r(x, t)e_2 + s(x, t)e_3 + t(x, t)e_4 + v(x, t)e_5 + w(x, t)e_6 + z(x, t)e_7 \in \mathbb{O}$ the symmetric octonionic KdV equation, in this form of (4.6) becomes a set of eight coupled equations

$$\begin{aligned} p_t + 6pp_x - 6qq_x - 6rr_x - 6ss_x - 6tt_x - 6vv_x - 6ww_x - 6zz_x + p_{xxx} &= 0, \\ \chi_t + 6\chi p_x + 6p\chi_x + \chi_{xxx} &= 0, \end{aligned} \quad (5.4)$$

with $\chi = q, r, s, t, v, w, z$. Setting any of four variables for χ to zero reduces (5.4) to the coupled set of equations corresponding to the symmetric quaternionic KdV equation (3.5) with constraints (3.7).

5.2. $\mathcal{PT}_{e_1 e_2 e_3 e_4 e_5 e_6 e_7}$ -symmetric N -soliton solutions

Using the representation (5.2) we proceed as in subsection 3.3 and consider the shifted solution (2.24) in the complex space $\mathbb{C}(o)$

$$u_{a_1 \ell + o \mathcal{O}, \alpha}(x, t) = p_{a_1, \mathcal{O}, \alpha}(x, t) - o q_{a_1, \mathcal{O}, \alpha}(x, t) \quad (5.5)$$

$$= p_{a_1, \mathcal{O}, \alpha}(x, t) \ell - \frac{1}{\mathcal{O}} q_{a_1, \mathcal{O}, \alpha}(x, t) \sum_{i=1}^7 a_i e_i \quad (5.6)$$

that solves the octonionic KdV equation (5.4). The solution in (5.5) is $\mathcal{PT}_{e_1 e_2 e_3 e_4 e_5 e_6 e_7}$ -symmetric. Once more, multi-soliton solutions can be constructed in analogy to the complex case $\mathbb{C}(i)$ treated in [13] by treating all functions in $\mathbb{C}(o)$ as explained in more detail at the end of section 4.

6. Conclusions

We have shown that the bicomplex, quaternionic, coquaternionic and octonionic versions of the KdV equation admit multi-soliton solutions. Using the standard folklore we assume that the existence of such type of solutions implies certain integrability of these equations, which we did not formally prove. The bicomplex versions, local and nonlocal, display a particularly rich structure with the two types of solutions found to exhibit very different types of qualitative behaviour. Especially interesting is the solution in the idempotent representation that decomposes a N -soliton into a $2N$ -solitonic structure. Each one-soliton constituent of the N -soliton has two contributions that even involve two

independent speed parameters. Unlike as for the real and complex solitons, where the degeneracy poses a nontrivial technical problem [11, 15], here these parameters can be trivially set to be equal.

For all noncommutative versions of the KdV equation, i.e. quaternionic, coquaternionic and octonionic, we found multi-soliton solutions based on complex representation in which the imaginary unit is built from specific combinations of the imaginary and hyperbolic units. Interestingly in all cases we observe that the \mathcal{PT} -symmetry needed to ensure that the newly defined imaginary unit can also be used as a \mathcal{PT} -symmetry imposes constraints that are equivalent to the constraints needed to obtain the symmetric KdV equation from the nonsymmetric one.

Naturally it would be interesting to extend the analysis presented here to other types of nonlinear integrable systems. A more challenging extension is to multi-complexify also the variables x and t which then also impacts on the definition of the derivatives with respect to these variables.

Acknowledgments

JC is supported by a City, University of London Research Fellowship.

References

- [1] M.J. Ablowitz and Z.H. Musslimani, Integrable nonlocal nonlinear Schrödinger equation, *Phys. Rev. Lett.*, **110** (2013), 064105 (5).
- [2] S.L. Adler, *Quaternionic quantum mechanics and quantum fields*, **88** Oxford University Press on Demand (1995).
- [3] B. Bagchi and A. Banerjee, Bicomplex hamiltonian systems in quantum mechanics, *J. of Phys. A: Math. and Theor.*, **48** (2015), 505201 (29).
- [4] A. Banerjee, On the quantum mechanics of bicomplex Hamiltonian system, *Ann. of Phys.*, **377** (2017), 493–505.
- [5] A. Banerjee, Bicomplex Harmonic and Isotonic Oscillators: The Excited States, *Advances in Applied Clifford Algebras*, **27** (2017), 2321–2332.
- [6] A. Banerjee and A. Biswas, Exact bound state solutions for the bicomplex Morse oscillator, in: *AIP Conference Proceedings*, **1975** (2018), 030001(10).
- [7] C.M. Bender, Making sense of non-Hermitian Hamiltonians, *Rept. Prog. Phys.*, **70** (2007), 947–1018.
- [8] C.M. Bender and S. Boettcher, Real Spectra in Non-Hermitian Hamiltonians Having PT Symmetry, *Phys. Rev. Lett.*, **80** (1998), 5243–5246.
- [9] D.C. Brody and E.-M. Graefe, On complexified mechanics and coquaternions, *J. of Phys. A: Math. and Theor.*, **44** (2011), 072001(9).
- [10] J. Cen, F. Correa, and A. Fring, Integrable nonlocal Hirota equations, *J. Math. Phys.*, **60** (2019), 081508(18).
- [11] J. Cen, F. Correa, and A. Fring, Degenerate multi-solitons in the sine-Gordon equation, *J. Phys. A: Math. Theor.*, **50** (2017), 435201(20).
- [12] J. Cen, F. Correa, and A. Fring, Time-delay and reality conditions for complex solitons, *J. of Math. Phys.*, **58** (2017), 032901(14).
- [13] J. Cen and A. Fring, Complex solitons with real energies, *J. Phys. A: Math. Theor.*, **49** (2016), 365202(15).
- [14] J. Cen and A. Fring, Asymptotic and scattering behaviour for degenerate multi-solitons in the Hirota equation, *Physica D: Nonlinear Phenomena*, **397** (2019), 17–24.
- [15] F. Correa and A. Fring, Regularized degenerate multi-solitons, *Journal of High Energy Physics*, **2016** (2016), 8 (15).
- [16] D. Dast, D. Haag, H. Cartarius, J. Main, and G. Wunner, Eigenvalue structure of a Bose–Einstein condensate in a-symmetric double well, *J. of Phys. A: Math. and Theor.*, **46** (2013), 375301(19).
- [17] C.M. Davenport, A commutative hypercomplex algebra with associated function theory, in: *Clifford algebras with numeric and symbolic computations*, (1996) 213–227.

- [18] D. Dizdarevic, D. Dast, D. Haag, J. Main, H. Cartarius, and G. Wunner, Cusp bifurcation in the eigenvalue spectrum of PT-symmetric Bose-Einstein condensates, *Phys. Rev. A*, **91** (2015), 033636(6).
- [19] D. Finkelstein, J.M. Jauch, S. Schiminovich, and D. Speiser, Foundations of quaternion quantum mechanics, *J. of Math. Phys.*, **3** (1962), 207–220.
- [20] P.R. Girard, The quaternion group and modern physics, *Euro. J. of Phys.*, **5** (1984), 25–32.
- [21] R. Gutöhrlein, H. Cartarius, J. Main, and G. Wunner, Bifurcations and exceptional points in a-symmetric dipolar Bose–Einstein condensate, *J. of Phys. A: Math. and Theor.*, **49** (2016), 485301(18).
- [22] M. Günaydin and F. Gürsey, Quark structure and octonions, *J. of Math. Phys.*, **14** (1973), 1651–1667.
- [23] S.Y. Lou, Alice-Bob systems, P_s - T_d - C principles and multi-soliton solutions, arXiv preprint *arXiv:1603.03975* (2016).
- [24] S.Y. Lou and F. Huang, Alice-Bob physics: coherent solutions of nonlocal KdV systems, *Scientific Reports*, **7** (2017), 869(11).
- [25] S.Y. Lou, Alice-Bob systems, P - T - \hat{C} symmetry invariant and symmetry breaking soliton solutions, *J. of Math. Phys.*, **59** (2018), 083507(20).
- [26] M.E. Luna-Elizarraras, M. Shapiro, D.C. Struppa, and A. Vajiac, Bicomplex numbers and their elementary functions, *Cubo (Temuco) A Mathematical Journal*, **14** (2012), 61–80.
- [27] K. Manikandan, S. Stalin, and M. Senthilvelan, Dynamical behaviour of solitons in a PT-invariant nonlocal nonlinear Schrödinger equation with distributed coefficients, *The Europ. Phys. J. B*, **91** (2018), 291(11).
- [28] A. Mostafazadeh, Pseudo-Hermitian Representation of Quantum Mechanics, *Int. J. Geom. Meth. Mod. Phys.*, **7** (2010), 1191–1306.
- [29] G.B. Price, *An introduction to multicomplex spaces and functions*, M. Dekker inc, New York (1991).
- [30] D. Rochon and M. Shapiro, On algebraic properties of bicomplex and hyperbolic numbers, *Anal. Univ. Oradea, Fasc. Math.*, **11** (2004), 110(28).
- [31] S. Sangwine, T. Ell, and N. Le Bihan, Fundamental representations and algebraic properties of biquaternions or complexified quaternions, *Advances in Applied Clifford Algebras*, **21** (2011), 607–636.
- [32] F.G. Scholtz, H.B. Geyer, and F. Hahne, Quasi-Hermitian Operators in Quantum Mechanics and the Variational Principle, *Ann. Phys.*, **213** (1992), 74–101.
- [33] S. Stalin, M. Senthilvelan, and M. Lakshmanan, Nonstandard bilinearization of PT-invariant nonlocal nonlinear Schrödinger equation: Bright soliton solutions, *Phys. Lett. A*, **381** (2017), 2380–2385.
- [34] G. Sobczyk, The hyperbolic number plane, *The College Math. J.*, **26** (1995), 268–280 .
- [35] K.A. Theaker and R.A. Van Gorder, Multicomplex wave functions for linear and nonlinear Schrödinger equations, *Advances in Applied Clifford Algebras*, **27** (2017), 1857–1879.
- [36] S. Ulrych, Relativistic quantum physics with hyperbolic numbers, *Phys. Lett. B*, **625** (2005), 313–323.
- [37] A. Vourdas, Quantum systems with finite Hilbert space: Galois fields in quantum mechanics, *J. of Phys. A: Math. and Theor.*, **40** (2007), R285–R331.
- [38] Y. Xuegang, Hyperbolic Hilbert Space, *Advances in Applied Clifford Algebras*, **10** (2000), 49(12).