



Journal of Nonlinear Mathematical Physics

ISSN (Online): 1776-0852

ISSN (Print): 1402-9251

Journal Home Page: <https://www.atlantis-press.com/journals/jnmp>

Ermakov-Painlevé II Reduction in Cold Plasma Physics. Application of a Bäcklund Transformation

Colin Rogers, Peter A. Clarkson

To cite this article: Colin Rogers, Peter A. Clarkson (2018) Ermakov-Painlevé II Reduction in Cold Plasma Physics. Application of a Bäcklund Transformation, Journal of Nonlinear Mathematical Physics 25:2, 247–261, DOI:

<https://doi.org/10.1080/14029251.2018.1452672>

To link to this article: <https://doi.org/10.1080/14029251.2018.1452672>

Published online: 04 January 2021

Ermakov-Painlevé II Reduction in Cold Plasma Physics. Application of a Bäcklund Transformation

Colin Rogers

*School of Mathematics and Statistics, The University of New South Wales,
Sydney, NSW2052, Australia
c.rogers@unsw.edu.au*

Peter A. Clarkson

*School of Mathematics, Statistics & Actuarial Science,
University of Kent, Canterbury, CT2 7FS, UK
P.A.Clarkson@kent.ac.uk*

Received 16 September 2017

Accepted 9 December 2017

A class of symmetry transformations of a type originally introduced in a nonlinear optics context is used here to isolate an integrable Ermakov-Painlevé II reduction of a resonant NLS equation which encapsulates a nonlinear system in cold plasma physics descriptive of the uni-axial propagation of magneto-acoustic waves. A Bäcklund transformation is employed in the iterative generation of novel classes of solutions to the cold plasma system which involve either Yablonski-Vorob'ev polynomials or classical Airy functions.

1. Introduction

The extensive connections between plasma physics, magnetohydrodynamics and canonical nonlinear integrable equations of modern soliton theory are well-documented. These links originated with pioneering work by Washimi and Taniuti [72] who employed reductive perturbation techniques to derive the Korteweg-de Vries (KdV) equation in the analysis of the propagation of ion-acoustic waves in a plasma. In contemporary work, Berezhin and Karpman [7, 8] independently derived the KdV equation in studies, both numerical and analytic, of large amplitude disturbances in plasmas and other dispersive media. Reviews of the role of the KdV equation in plasma physics and of the application of the reductive perturbation method to the study of hydrodynamic waves in cold plasma have been presented in [24] and [26] respectively. The integrable mKdV equation, related to the KdV equation by a Miura transformation, likewise arises in the analysis of the propagation of nonlinear Alfvén waves in cold collisionless plasma [27]. In addition, the nonlinear Schrödinger equation has an extensive literature concerned with its derivation in plasma physics in a variety of contexts. Thus, in particular, its occurrence in the analysis of the propagation of Langmuir waves in plasma has been described in [18, 23, 70]. The text [73] provides an account of perturbation methods adduced in the derivation of the NLS equation in plasma physics, notably with regard to incorporation of Landau damping phenomena.

An integrable Heisenberg spin equation associated with the NLS equation has been derived in spatial gasdynamics and magneto-hydrostatics in [45, 46] via geometric methods originally applied in a magnetohydrodynamic context in [47]. This purely geometric formalism as opposed to a reductive perturbation approach may be shown to lead to a classical integrable da Rios system reduction

in complex-lamellar magnetohydrodynamics [48]. It has also been applied not only to construct the auto-Bäcklund transformation for the auto-Bäcklund transformation for the NLS equation in a purely geometric manner [49, 50] but also to obtain a novel integrable Pohlmeyer-Lund-Regge reduction in magnetohydrodynamics [51, 66]. The combined action of Bäcklund and reciprocal-type transformations has been used, via an integrable sinh-Gordon reduction, to construct periodic solutions of breather-type in super-Alfvénic magnetogasdynamics in [52]. Invariance under multi-parameter reciprocal transformations has been established in two-dimensional orthogonal magnetogasdynamics in [53].

In [54, 55] elliptic vortex solutions in magnetogasdynamics have been obtained via symmetry reduction to an integrable Ermakov-Ray-Reid system with underlying Hamiltonian structure. Ermakov-Ray-Reid systems have diverse physical applications, in particular, in nonlinear optics and rotating shallow water theory (see e.g. [56–58]). Moreover, in [59], a 2 + 1-dimensional non-isothermal magnetogasdynamic version of a gas cloud system with origin in work of Dyson [15] was shown to admit symmetry reduction to an eight dimensional nonlinear dynamical subsystem with underlying Hamiltonian-Ermakov structure. A Lax pair representation was constructed for this integrable subsystem.

Here, a class of wave packet representations of a type originally introduced in a nonlinear optics context in [20] is used to isolate a Ermakov-Painlevé II symmetry reduction of a resonant NLS equation which encapsulates a 1 + 1-dimensional cold plasma physics system. The latter describes the propagation of uniaxial long magnetoacoustic waves in a cold collisionless plasma subject to a transverse magnetic field. Iterative application of a Bäcklund transformation is used to generate novel classes of exact solutions of the cold plasma system in terms of either Yablonski-Vorob’ev polynomials or classical Airy functions.

2. The Two-Component Cold Plasma System. A Resonant NLS Encapsulation

The dynamics of two-component cold collisionless plasma in the presence of an external magnetic field \mathbf{B} is governed by the nonlinear system of equations [3, 28].

$$m_i[\partial/\partial t + \mathbf{v}_i \nabla] \mathbf{v}_i = e[\mathbf{E} + (\mathbf{v}_i \times \mathbf{B})], \quad (2.1a)$$

$$m_e[\partial/\partial t + \mathbf{v}_e \nabla] \mathbf{v}_e = -e[\mathbf{E} + (\mathbf{v}_e \times \mathbf{B})], \quad (2.1b)$$

$$\frac{\partial n_i}{\partial t} + \nabla \cdot (n_i \mathbf{v}_i) = 0 \quad (2.1c)$$

$$\frac{\partial n_e}{\partial t} + \nabla \cdot (n_e \mathbf{v}_e) = 0, \quad (2.1d)$$

$$\text{curl } \mathbf{B} = e\mu_0(n_i \mathbf{v}_i - n_e \mathbf{v}_e), \quad (2.1e)$$

$$\frac{\partial \mathbf{B}}{\partial t} = -\text{curl } \mathbf{E}, \quad (2.1f)$$

$$\nabla \cdot \mathbf{B} = 0 \quad (2.1g)$$

where $m_i, m_e, \mathbf{v}_i, \mathbf{v}_e, n_i, n_e$ denote, in turn, masses, velocities and concentrations of ions and electrons respectively. \mathbf{E} is the electric field, \mathbf{B} is the magnetic field, e is the electric charge and μ_0 is the magnetic permeability. If the oscillation frequency is much smaller than the ion Langmuir frequency, then plasma quasi-neutrality is implied, that is, $n_i \approx n_e = n$.

On introduction of the mass density ρ and velocity \mathbf{u} according to

$$\rho = (m_i + m_e)n, \quad \mathbf{u} = \frac{m_i \mathbf{v}_i + m_e \mathbf{v}_e}{m_i + m_e},$$

then, on elimination of the electric field \mathbf{E} , if $m_e/m_i \ll 1$, in the case of uni-axial plasma propagation with transverse magnetic field \mathbf{B} , so that

$$\mathbf{u} = u(x,t)\mathbf{i}, \quad \mathbf{B} = B(x,t)\mathbf{k}, \quad (2.2)$$

the system (2.1) may be reduced to the form [33]

$$\frac{\partial \rho}{\partial t} + \frac{\partial}{\partial x}(\rho u) = 0, \quad (2.3a)$$

$$\rho \left(\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} \right) + B \frac{\partial B}{\partial x} = 0, \quad (2.3b)$$

$$\frac{\partial}{\partial t} \left[B - \frac{\partial}{\partial x} \left(\frac{1}{\rho} \frac{\partial B}{\partial x} \right) \right] + \frac{\partial}{\partial x} \left\{ u \left[B - \frac{\partial}{\partial x} \left(\frac{1}{\rho} \frac{\partial B}{\partial x} \right) \right] \right\} = 0. \quad (2.3c)$$

Introduction of the Lagrangian variable $X(x,t)$ via the continuity equation (2.3a) according to

$$dX = \rho dx - \rho u dt,$$

into (2.3c) shows that

$$B - \frac{\partial}{\partial x} \left(\frac{1}{\rho} \frac{\partial B}{\partial x} \right) = M(X)\rho. \quad (2.4)$$

The system consisting of (2.3a), (2.3b) augmented by (2.4) with $M(X) = 1$ was set down in a non-linear dispersive wave context by Whitham [74]. In the present context it describes the propagation of 1 + 1-dimensional nonlinear magnetoacoustic waves in a cold plasma subject to a transverse magnetic field.

Here, we consider a shallow water type approximation to the Whitham system consisting of the continuity and momentum equations (2.3a), (2.3b) together with

$$B - \frac{\partial}{\partial x} \left(\frac{1}{\rho} \frac{\partial B}{\partial x} \right) = \rho. \quad (2.5)$$

Thus, on re-scaling the space and time variables via $x \rightarrow \beta x$, $t \rightarrow \beta t$ and expansion of the expression for the magnetic field B in the parameter β^2 according to [33]

$$B = \rho + \beta^2 \mathcal{B}(\rho, \rho_x, \rho_{xx}, \dots) + \mathcal{O}(\beta^4), \quad (2.6)$$

insertion into (2.5) yields

$$\mathcal{B} = \frac{\partial}{\partial x} \left(\frac{1}{\rho} \frac{\partial \rho}{\partial x} \right). \quad (2.7)$$

Thus, to $\mathcal{O}(\beta^2)$, the following nonlinear system results [33]:

$$\frac{\partial \rho}{\partial t} + \frac{\partial}{\partial x}(\rho u) = 0, \quad (2.8a)$$

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} + \beta^2 \frac{\partial}{\partial x} \left[\frac{1}{\rho} \frac{\partial^2 \rho}{\partial x^2} - \frac{1}{2} \left(\frac{1}{\rho} \frac{\partial \rho}{\partial x} \right)^2 + \frac{\rho}{\beta^2} \right]. \quad (2.8b)$$

This describes the uniaxial propagation of long magneto-acoustic waves in a cold plasma with velocity magnitude u and magnetic field given by (2.2) together with (2.6)–(2.7).

On use of the one-dimensional version of a relation for the de Broglie-Bohm potential, namely

$$\frac{\nabla^2 \rho^{1/2}}{\rho^{1/2}} = \frac{\nabla^2 \rho}{\rho} - \frac{1}{4} \left(\frac{\nabla \rho}{\rho} \right)^2,$$

it seen that the cold plasma momentum equation in (2.8a) becomes

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} + \beta^2 \frac{\partial}{\partial x} \left[2 \frac{(\rho^{1/2})_{xx}}{\rho^{1/2}} + \frac{\rho}{\beta^2} \right] = 0. \quad (2.9)$$

Introduction of the velocity potential S according to $u = -2\partial S/\partial x$ into the continuity equation (2.8a) and momentum equation (2.9) in turn, gives

$$\frac{\partial \rho}{\partial t} - 2 \frac{\partial}{\partial x} \left(\rho \frac{\partial S}{\partial x} \right) = 0,$$

together with the Bernoulli integral

$$-\frac{\partial S}{\partial t} + \left(\frac{\partial S}{\partial x} \right)^2 + \frac{\rho}{2} + \frac{\beta^2}{\rho^{1/2}} \frac{\partial^2}{\partial x^2} (\rho^{1/2}) = T(t).$$

In the latter, $T(t)$ may be absorbed into the potential S and so may be set zero without loss of generality.

If the classical Madelung transformation is now introduced according to [38]

$$\Psi = \rho^{1/2} e^{-iS}$$

then it is seen that the cold plasma system (2.8a) may be encapsulated in the resonant NLS equation [43, 44]

$$i \frac{\partial \Psi}{\partial t} + \frac{\partial^2 \Psi}{\partial x^2} - \frac{1}{2} |\Psi|^2 \Psi = (1 + \beta^2) \frac{\Psi}{|\Psi|} \frac{\partial^2 |\Psi|}{\partial x^2}, \quad (2.10)$$

which incorporates a de Broglie-Bohm potential term $\frac{1}{|\Psi|} \frac{\partial^2}{\partial x^2} |\Psi|$. It is recalled that, if $s < 1$ then the canonical 1 + 1-dimensional resonant NLS equation

$$i \frac{\partial \Psi}{\partial t} + \frac{\partial^2 \Psi}{\partial x^2} + v |\Psi|^2 \Psi = \frac{s}{|\Psi|} \frac{\partial^2 |\Psi|}{\partial x^2},$$

can be transformed to the standard integrable cubic NLS equation with the de Broglie-Bohm term removed [43, 44, 60]. However, if as in the present cold plasma case $s = 1 + \beta^2 > 1$ this reduction to remove the de Broglie Bohm term is not available. Then, as shown in [33] the resonant NLS equation (2.10) which encapsulates the cold plasma system is equivalent to a canonical two-component system contained in the AKNS hierarchy of solitonic systems amenable to the inverse scattering transform [1, 2]. Bäcklund-Darboux transformations and concomitant nonlinear superposition principles for the resonant NLS equation and hence for the cold plasma system which it encapsulates have been constructed in [33]. This resonant NLS equation importantly, unlike its standard integrable NLS counterpart can admit solitonic fusion or fission phenomena [44]. Here, wave packet

solutions of the resonant NLS equation (2.10) are generated via a symmetry reduction to a prototype integrable Ermakov-Painlevé II equation [4, 61]. The iterated application of a Bäcklund transformation then allows the construction of novel classes of exact solutions to the cold plasma system in terms of either Yablonski-Vorob'ev polynomials or classical Airy functions.

3. Ermakov-Painlevé II Symmetry Reduction

Here, symmetry reduction of the resonant NLS equation (2.10) is sought under the wave packet ansatz

$$\Psi = \rho^{1/2} e^{-iS} = [\phi(\xi) + i\psi(\xi)] e^{i\eta}, \quad (3.1a)$$

$$\xi = \alpha t + \beta t^2 + \gamma x, \quad (3.1b)$$

$$\eta = \gamma t^3 + \delta t^2 + \varepsilon \gamma t x + \zeta t + \lambda x, \quad (3.1c)$$

with $\alpha, \beta, \gamma, \delta, \varepsilon, \zeta$ and λ arbitrary constants. Insertion of the latter into (2.10) produces the coupled nonlinear system

$$\gamma^2 \frac{d^2 \phi}{d\xi^2} - \frac{d\psi}{d\xi} [2(\beta + \varepsilon \gamma^2)t + \alpha + 2\lambda \gamma] - \Delta \phi = 0, \quad (3.2a)$$

$$\gamma^2 \frac{d^2 \psi}{d\xi^2} + \frac{d\phi}{d\xi} [2(\beta + \varepsilon \gamma^2)t + \alpha + 2\lambda \gamma] - \Delta \psi = 0, \quad (3.2b)$$

where

$$\Delta = 3\gamma t^2 + 2\delta t + \varepsilon \gamma x + \zeta + (\varepsilon \gamma t + \lambda)^2 + \frac{1}{2} |\Psi|^2 + \frac{s\gamma^2}{|\Psi|} \frac{d^2 |\Psi|}{d\xi^2}. \quad (3.3)$$

and $s = 1 + \beta^2 > 1$.

The relations (3.2) combine to show that

$$\gamma^2 \left(\psi \frac{d^2 \phi}{d\xi^2} - \phi \frac{d^2 \psi}{d\xi^2} \right) - \left(\phi \frac{d\phi}{d\xi} + \psi \frac{d\psi}{d\xi} \right) [2(\beta + \varepsilon \gamma^2)t + \alpha + 2\lambda \gamma] = 0, \quad (3.4)$$

whence, it is required that

$$\beta + \varepsilon \gamma^2 = 0,$$

in which case, (3.4) admits the integral

$$\gamma^2 \left(\psi \frac{d\phi}{d\xi} - \phi \frac{d\psi}{d\xi} \right) - \frac{1}{2} (\alpha + 2\lambda \gamma) |\Psi|^2 = \mathcal{I}, \quad (3.5)$$

where \mathcal{I} is an arbitrary constant.

It is seen that (3.3) may be written as, if $\beta \neq 0$

$$\begin{aligned} \Delta &= \gamma\beta^{-1}(\varepsilon^2\gamma + 3)(\xi - \alpha t - \gamma x) + 2(\delta + \varepsilon\gamma\lambda)t + \varepsilon\gamma x + \zeta + \lambda^2 \\ &\quad + \frac{1}{2}|\Psi|^2 + \frac{s\gamma^2}{|\Psi|} \frac{d^2|\Psi|}{d\xi^2} \\ &= \varepsilon\xi + \zeta + \lambda^2 + \frac{1}{2}|\Psi|^2 + \frac{s\gamma^2}{|\Psi|} \frac{d^2|\Psi|}{d\xi^2}, \end{aligned} \tag{3.6}$$

on setting

$$\alpha\varepsilon = 2(\delta + \varepsilon\gamma\lambda), \quad \beta\varepsilon = \gamma(3 + \varepsilon^2\gamma).$$

Moreover, (3.2) show that

$$\gamma^2 \left(\phi \frac{d^2\phi}{d\xi^2} + \psi \frac{d^2\psi}{d\xi^2} \right) + \left(\psi \frac{d\phi}{d\xi} - \phi \frac{d\psi}{d\xi} \right) (\alpha + 2\lambda\gamma) - \Delta|\Psi|^2 = 0,$$

whence, on use of the relations

$$\left[\phi \frac{d^2\phi}{d\xi^2} + \psi \frac{d^2\psi}{d\xi^2} + \left(\frac{d\phi}{d\xi} \right)^2 + \left(\frac{d\psi}{d\xi} \right)^2 \right] (\phi^2 + \psi^2) - \left(\phi \frac{d\phi}{d\xi} + \psi \frac{d\psi}{d\xi} \right)^2 = \frac{d^2|\Psi|}{d\xi^2} |\Psi|^3,$$

and

$$(\phi^2 + \psi^2) \left[\left(\frac{d\phi}{d\xi} \right)^2 + \left(\frac{d\psi}{d\xi} \right)^2 \right] - \left(\psi \frac{d\phi}{d\xi} - \phi \frac{d\psi}{d\xi} \right)^2 = \left(\phi \frac{d\phi}{d\xi} + \psi \frac{d\psi}{d\xi} \right)^2,$$

it is seen that

$$\gamma^2 \left[\frac{d^2|\Psi|}{d\xi^2} |\Psi|^3 - \left(\psi \frac{d\phi}{d\xi} - \phi \frac{d\psi}{d\xi} \right)^2 \right] + (\alpha + 2\lambda\gamma) \left(\psi \frac{d\phi}{d\xi} - \phi \frac{d\psi}{d\xi} \right) |\Psi|^2 - \Delta|\Psi|^4 = 0.$$

The latter, by virtue of the integral of motion (3.5) and the expression (3.6) for Δ now produces a hybrid Ermakov-Painlevé equation in the amplitude $|\Psi|$ namely

$$|\Psi|_{\xi\xi} + (c_1 + c_2\xi)|\Psi| + c_3|\Psi|^3 = \frac{\mathcal{I}^2}{(1-s)\gamma^4|\Psi|^3}, \tag{3.7}$$

where the constants c_1, c_2 and c_3 are given by

$$c_1 = \frac{1}{(1-s)\gamma^4} \left[\left(\alpha - \frac{\delta}{\varepsilon} \right)^2 - \gamma^2(\zeta + \lambda^2) \right], \quad c_2 = \frac{\varepsilon}{(s-1)\gamma^2}, \quad c_3 = \frac{1}{2(s-1)\gamma^2}.$$

The type of nonlinear equation (3.7) has been previously shown to arise in connection with a pair of three-ion cases in the classical Nernst-Planck electrodiffusion system [4]. Here, $|\Psi| = \rho^{1/2}$ and

in terms of the density ρ it is seen that (3.7) becomes

$$\frac{d^2\rho}{d\xi^2} - \frac{1}{2\rho} \left(\frac{d\rho}{d\xi}\right)^2 + 2[c_1 + c_2\xi]\rho + 2c_3\rho^2 + \frac{2\mathcal{I}^2}{(s-1)\gamma^4\rho} = 0,$$

which with the admissible specialisations,

$$c_1 = 0, \quad c_2 = \frac{1}{2}, \quad c_3 = 1, \quad \frac{4\mathcal{I}^2}{(s-1)\gamma^4} = (\alpha + \frac{1}{2})^2, \tag{3.8}$$

becomes the canonical integrable Painlevé XXXIV (P₃₄) equation in $\rho > 0$, namely

$$\frac{d^2\rho}{d\xi^2} = \frac{1}{2\rho} \left(\frac{d\rho}{d\xi}\right)^2 + 2\rho^2 - \xi\rho - \frac{(\alpha + \frac{1}{2})^2}{2\rho}, \tag{3.9}$$

cf. Gromak [21]. It is emphasised that the relation in (3.8) involving the Painlevé parameter α , necessarily requires that $s > 1$ as indeed is the case for the present cold plasma system incapsulated in a nonlinear resonant NLS equation. The regions of positivity of classes of solutions of P₃₄ (3.9) as generated by the iterated action of a Bäcklund transformation have been investigated in the context of boundary value problems in two-ion electrodiffusion in [5]. Therein, the ion concentrations c_{\pm} , which are necessarily positive, are governed by solutions of P₃₄ (3.9).

With a positive solution $\rho = |\Psi|^2 = \phi^2 + \psi^2$ of P₃₄ (3.9) to hand, the ratio ψ/ϕ is determined by the integral of motion (3.5). The latter yields

$$-\gamma^2 \frac{d}{d\xi} \left[\tan^{-1} \left(\frac{\psi}{\phi} \right) \right] - \frac{1}{2}(\alpha + 2\lambda\gamma) = \frac{\mathcal{I}}{|\Psi|^2},$$

so that, on integration,

$$\gamma^2 \tan^{-1} \left(\frac{\psi}{\phi} \right) + \frac{1}{2}(\alpha + 2\lambda\gamma)\xi + \mathcal{I} \int \frac{d\xi}{\rho} = K,$$

where K is an arbitrary constant. Thus, with $\Lambda = \psi/\phi$, it is seen that the ϕ, ψ in the original wave packet representation (3.1) are given by the relations

$$\phi = \pm \frac{|\Psi|}{\sqrt{1+\Lambda^2}}, \quad \psi = \pm \frac{\Lambda|\Psi|}{\sqrt{1+\Lambda^2}}.$$

The velocity magnitude in the cold plasma system (2.8a) is given by $u = \Phi_x$ where Φ is the potential given by

$$\Phi = 2 \tan^{-1} \left(\frac{\psi + \phi \tan \eta}{\phi - \psi \tan \eta} \right) = 2 \left[\tan^{-1} \left(\frac{\psi}{\phi} \right) + \eta \right],$$

on use of the identity

$$\tan^{-1} \left(\frac{x+y}{1-xy} \right) \equiv \tan^{-1} x + \tan^{-1} y.$$

Thus,

$$\begin{aligned} u &= 2 \left[\gamma \frac{\partial}{\partial \xi} + (\varepsilon\gamma + \lambda) \frac{\partial}{\partial \eta} \right] \left[-\frac{1}{2\gamma^2}(\alpha + 2\lambda\gamma)\xi - \frac{\mathcal{I}}{\gamma^2} \int \frac{d\xi}{\rho(\xi)} + \eta \right] \\ &= -\frac{2\mathcal{I}}{\gamma\rho(\xi)} + 2\varepsilon\gamma - \frac{\alpha}{\gamma}. \end{aligned}$$

while the term \mathcal{B} in the expansion (2.6) for the magnetic field B is given by

$$\mathcal{B} = \gamma^2 \frac{d^2}{d\xi^2} \ln \rho(\xi).$$

where the density $\rho(\xi)$ is governed by P_{34} (3.9).

In the sequel, a link between the Ermakov-Painlevé II equation for the density $\rho(\xi)$ and the Painlevé II (P_{II}) equation

$$\frac{d^2 q}{d\xi^2} = 2q^3 + \xi q + \alpha, \quad (3.10)$$

with α a parameter, via P_{34} (3.9) is exploited to construct novel classes of wave packet solutions of the cold plasma system in terms of Yablonski-Vorob'ev polynomials or classical Airy functions via the iterated application of the well-known Bäcklund transformation for P_{II} [19,37]; see also [10,22]. These solutions may, in turn, be used as seed solutions in the application of the Bäcklund-Darboux transformations as established for the resonant NLS equation in [33].

4. Iterative Action of a Bäcklund Transformation

Bäcklund transformations have established applications in continuum mechanics and notably in modern soliton theory [50,62]. These have roots, in turn, in work of Loewner [34,35] on model laws in gasdynamics and Seeger *et al.* [69] on crystal dislocations. Thus, Loewner applied novel matrix Bäcklund transformations to construct multi-parameter gas laws for which the classical hodograph system may be reduced to appropriate canonical forms in subsonic, transonic and supersonic flow régimes. Seeger *et al.* [69], on the other hand, in the context of Frenkel and Kontorova's dislocation theory isolated the nonlinear interaction of what they termed 'eigenmotions' via Bianchi's classical permutability theorem as derived via the auto-Bäcklund transformation for the sine Gordon equation of pseudo-spherical surface theory. In particular, the interaction process of what, in soliton theory came to be called a breather and kink was described analytically by means of this nonlinear superposition principle. Importantly, Lamb [32] later exploited this permutability theorem to predict experimentally observed decomposition of ultrashort optical pulses in a resonant medium.

The areas of application of Bäcklund transformations in nonlinear continuum mechanics and modern soliton theory were brought together in [29, 30] where it was established that a re-interpretation and extension of the class of matrix infinitesimal Bäcklund transformations introduced by Loewner in a gasdynamics setting provide a linear representation for a master 2 + 1-dimensional soliton system. Basic reductions of this system lead, in particular to novel integrable 2 + 1-dimensional versions of the principal chiral fields model, Toda lattice system and, notably, of the classical sine Gordon equation [9, 11, 30, 31, 36, 40, 41, 67, 68]. It is remarked that the paper of Loewner [34] contains, 'mutatis mutandis' a linear representation for the 1 + 1-dimensional sine Gordon equation which is gauge equivalent to that later obtained in the celebrated AKNS system [1].

The preceding attests to the seminal role that Bäcklund transformations have played in modern soliton theory. Likewise the classical Painlevé equations and the Bäcklund transformations they admit arise naturally in the study of integrable solitonic systems (see e.g. [6, 10, 16, 21, 22, 39] and work cited therein). Here, our concern is with the application of a Bäcklund transformation for P_{II} (3.10) to the present symmetry reduction of the cold plasma system encapsulated in the resonant NLS equation (2.10).

It turns out that, remarkably, all known exact solutions of P_{II} (3.10) and hence also of P_{34} (3.9) may be generated via iteration of a Bäcklund transformation of (3.10) due to Gambier [19] and later Lukashevich [37].

The well-known relationship between solutions of P_{II} (3.10) and solutions of P_{34} (3.9) is derived via the Hamiltonian system

$$\frac{dq}{d\xi} = \frac{\partial \mathcal{H}_{II}}{\partial \rho}, \quad \frac{d\rho}{d\xi} = -\frac{\partial \mathcal{H}_{II}}{\partial q},$$

where the Hamiltonian $\mathcal{H}_{II}(\rho, q, \xi; \alpha)$ is given by

$$\mathcal{H}_{II}(\rho, q, \xi; \alpha) = \frac{1}{2}\rho^2 - (q^2 + \frac{1}{2}\xi)\rho - (\alpha + \frac{1}{2})q,$$

which yield the nonlinear system

$$\frac{dq}{d\xi} = \rho - q^2 - \frac{1}{2}\xi, \quad \frac{d\rho}{d\xi} = 2q\rho + \alpha + \frac{1}{2}, \tag{4.1}$$

see [25, 42]. Elimination of ρ and q successively in (4.1) yields P_{II} (3.10) and P_{34} (3.9).

If $q_\alpha(\xi) = q(\xi; \alpha)$ is a solution of P_{II} (3.10) with parameter α , then

$$q_{\alpha+1}(\xi) = -q_\alpha(\xi) - \frac{2\alpha + 1}{2q'_\alpha(\xi) + 2q_\alpha^2(\xi) + \xi}, \tag{4.2a}$$

$$q_{\alpha-1}(\xi) = -q_\alpha(\xi) - \frac{2\alpha - 1}{2q'_\alpha(\xi) - 2q_\alpha^2(\xi) + \xi}, \tag{4.2b}$$

with $' \equiv d/d\xi$, which are the Bäcklund transformations of P_{II} (3.10) [19, 37]. If $\rho_\alpha(\xi) = \rho(\xi; \alpha)$ is a solution of P_{34} (3.9) with parameter α , then from (4.1) we have

$$\rho_\alpha = \frac{dq_\alpha}{d\xi} + q_\alpha^2 + \frac{1}{2}\xi, \quad q_\alpha = \frac{1}{2\rho_\alpha} \left(\frac{d\rho_\alpha}{d\xi} - \alpha - \frac{1}{2} \right). \tag{4.3}$$

Consequently, from (4.2) and (4.3) we obtain the Bäcklund transformations of P_{34} (3.9) given by

$$\rho_{\alpha+1} = z - \rho_\alpha + \frac{1}{2\rho_\alpha^2} \left(\frac{d\rho_\alpha}{d\xi} + \alpha + \frac{1}{2} \right)^2, \tag{4.4a}$$

$$\rho_{\alpha-1} = z - \rho_\alpha + \frac{1}{2\rho_\alpha^2} \left(\frac{d\rho_\alpha}{d\xi} - \alpha - \frac{1}{2} \right)^2, \tag{4.4b}$$

see also [17, 42]

In the present plasma physics context, in particular, this produces an extensive class of exact rational solutions for the density ρ governed by P_{34} (3.9) given by

$$\rho = \rho_+(\xi; n) = \frac{Q_{n+1}(\xi)Q_{n-1}(\xi)}{2Q_n^2(\xi)}, \quad n \in \mathbb{N}, \tag{4.5}$$

with $\rho_+(\xi; 0) = \frac{1}{2}\xi$, corresponding to the parameters $\alpha = n$, where the $Q_n(\xi)$ are the *Yablonskii-Vorob'ev polynomials* determined by the quadratic recurrence relations [71, 75]

$$Q_{n+1}Q_{n-1} = \xi Q_n^2 + 4 \left[\left(\frac{dQ_n}{d\xi} \right)^2 - Q_n \frac{d^2 Q_n}{d\xi^2} \right],$$

with $Q_{-1}(\xi) = Q_0(\xi) = 1$. The rational solutions of P_{34} (3.9) can also be expressed in terms of determinants. Let $p_k(\xi)$ be the polynomial defined by

$$\sum_{k=0}^{\infty} p_k(\xi)\lambda^k = \exp\left(\xi\lambda - \frac{4}{3}\lambda^3\right), \quad (4.6)$$

and $\tau_n(\xi)$ be the Wronskian

$$\tau_n(\xi) = \mathcal{W}(p_1(\xi), p_3(\xi), \dots, p_{2n-1}(\xi)) \equiv \begin{vmatrix} p_1(\xi) & p_3(\xi) & \cdots & p_{2n-1}(\xi) \\ p_1'(\xi) & p_3'(\xi) & \cdots & p_{2n-1}'(\xi) \\ \vdots & \vdots & \ddots & \vdots \\ p_1^{(n-1)}(\xi) & p_3^{(n-1)}(\xi) & \cdots & p_{2n-1}^{(n-1)}(\xi) \end{vmatrix}, \quad (4.7)$$

for $n \geq 1$. Then the rational solution

$$\rho_+(\xi; n) = \frac{1}{2}\xi - 2\frac{d^2}{d\xi^2} \ln \tau_n(\xi),$$

satisfies P_{34} (3.9) with $\alpha = n$.

Iterated action of the Bäcklund transformations (4.4) for P_{34} (3.9) on the seed Airy-type solution with parameter $\alpha = \frac{1}{2}$ generates a class of exact solutions of P_{34} (3.9) given by

$$\rho\left(\xi; n - \frac{1}{2}\right) = \frac{u_{n-1}(\xi)u_{n+1}(\xi)}{2u_n^2(\xi)}, \quad n \in \mathbb{N}, \quad (4.8)$$

for the parameter $\alpha = n - \frac{1}{2}$ and where the sequence $\{u_n\xi\}$, for $n \geq 0$, is determined by the Toda recurrence relation

$$u_{n+1}u_{n-1} = 4 \left[\left(\frac{du_n}{d\xi}\right)^2 - u_n \frac{d^2u_n}{d\xi^2} \right] \quad (4.9)$$

with initial values $u_0(\xi) = 1$ and $u_1(\xi) = \varphi(\xi)$, where $\varphi(\xi)$ is governed by the classical Airy equation

$$\frac{d^2\varphi}{d\xi^2} + \frac{1}{2}\xi\varphi = 0, \quad (4.10)$$

i.e.

$$\varphi(\xi) = a\text{Ai}(z) + b\text{Bi}(z), \quad z = -2^{-1/3}\xi, \quad (4.11)$$

with $\text{Ai}(z)$ and $\text{Bi}(z)$ the Airy functions and a, b arbitrary parameters. The Airy-type solutions of P_{34} (3.9) can also be expressed in terms of determinants. Suppose that $\Phi_n(\xi)$ is the Hankel $n \times n$

determinant

$$\Phi_n(\xi) = \left[\frac{d^{j+k}}{d\xi^{j+k}} \varphi(\xi) \right]_{j,k=0}^{n-1}, \quad n \geq 1, \quad (4.12)$$

with $\varphi(\xi)$ given by (4.11) and $\Phi_0(\xi) = 1$, then for $n \geq 1$,

$$p(\xi; n - \frac{1}{2}) = -2 \frac{d^2}{d\xi^2} \ln \Phi_n(\xi),$$

satisfies P₃₄ (3.9) with $\alpha = n - \frac{1}{2}$, cf. [12].

In previous work in [5], solutions in terms of P₃₄ in a two-ion electro-diffusion context have been used to investigate certain boundary value problems. There the positivity requirement on the solutions arises via such a constraint on the ion concentrations. Here, it arises due to the positivity constraint on the density. The rational solutions of P₃₄ are not positive for all ξ . However, positivity may be established in certain regions $\xi > c$, a constant. Thus, the rational solutions of P₃₄ are given by

$$p[k] = \frac{1}{2}\xi - 2 \frac{d^2}{d\xi^2} \ln P_{k-1}(\xi), \quad p[-1] = \frac{1}{2}\xi$$

where the P_k are the Yablonski-Vorob'ev polynomials, cf. [13, 14]. If ξ_{k-1} denotes the largest real zero of the polynomial P_{k-1} for each $k \in \mathbb{N}$ then the constraint $\xi_{k-1} < c$ ensures positivity in the region $\xi > c$. The positivity property is illustrated graphically for $k = 0, 1, 2$ in [5].

The Airy-type solutions of P_{II} admit the representation [5, 12]

$$Y[k - \frac{1}{2}] = \frac{d}{d\xi} \ln \frac{\psi_{k-2}}{\psi_{k-1}} - \Phi,$$

wherein the sequence $\{\psi_k\}_{k \geq -1}$ is given by the recurrence relation

$$\psi_{k+1} \psi_{k-1} = 4 \left[\left(\frac{d\psi_k}{d\xi} \right)^2 - \psi_k \frac{d^2\psi_k}{d\xi^2} + \frac{1}{2}(k+1)(2\Phi^2 + \xi)\psi_k^2 \right]$$

together with the initial values $\psi_{-1} = 1, \psi_0 = 1$. In the above $\Phi = (\ln \phi)'$ where ϕ is, in general, given by (4.11). The corresponding Airy-type solutions of P₃₄ are determined by [5, 12]

$$p[k - \frac{1}{2}] = k(2\Phi^2 + \xi) - 2 \frac{d^2}{d\xi^2} \ln \psi_{k-1}.$$

If one proceeds with the case $b = 0$ in (4.11), i.e.

$$\phi(\xi) = \text{Ai}(-2^{-1/3}\xi)$$

where we have set $a = 1$, without loss of generality, then it was established in [5, 12] that, for any k , there exists a ξ_k such that $p[k - \frac{1}{2}]$ is non-singular and positive on $\xi < \xi_k$. This positivity property is depicted graphically for $k = 1, 2, \dots, 6$ in [12].

5. Conclusion

The aim of the present work has been to exploit an integrable Ermakov-Painlevé II reduction of a resonant NLS encapsulation of a nonlinear cold plasma system to generate classes of similarity solutions in terms of either Yablonskii-Vorob'ev polynomials or classical Airy functions. It is remarked that classes of similarity solutions to the related P_{II} equation have recently been shown to provide exact solutions to nonlinear moving boundary problems for certain solitonic equations [63–65] while similarity solutions in magneto-gasdynamics descriptive of plasma columns confined by moving boundaries have been isolated in [55].

References

- [1] M.J. Ablowitz, D.J. Kaup, A.C. Newell and H. Segur, Nonlinear evolution equations of physical significance, *Phys. Rev. Lett.* **31** (1973) 125–127.
- [2] M.J. Ablowitz and P.A. Clarkson, *Solitons, Nonlinear Evolution Equations and Inverse Scattering* (LMS Lecture Notes on Mathematics, vol. **149**), Cambridge University Press, Cambridge (1991).
- [3] A.I. Akhiezer, I.A. Akhiezer, R.V. Polovin, A.G. Sitenko and K.N. Stepanov, *Plasma Electrodynamics. Volume 1: Linear theory. Volume 2: Non-linear theory and fluctuations* (International Series of Monographs in Natural Philosophy, vols. **68, 80**) Pergamon Press, Oxford (1975).
- [4] P. Amster and C. Rogers, On a Ermakov-Painlevé II reduction in three-ion electrodiffusion. A Dirichlet boundary value problem, *Discrete and Continuous Dynamical Systems* **35** (2015) 3277–3292.
- [5] L.K. Bass, J. Nimmo, C. Rogers and W.K. Schief, Electrical structures of interfaces. A Painlevé II model, *Proc. Roy. Soc. Lond. A* **466** (2010) 2117–2136.
- [6] A.P. Bassom, P.A. Clarkson and A. C. Hicks, Bäcklund transformations and solution hierarchies for the fourth Painlevé equation, *Stud. Appl. Math.* **95** (1995) 1–71.
- [7] Y.A. Berezin and V.I. Karpman, Theory of non-stationary finite amplitude waves in low density plasma, *Soviet Phys. JETP* **19** (1974) 1265–1271.
- [8] Y.A. Berezin and V.I. Karpman, Nonlinear evolution of disturbances in plasmas and other dispersive media, *Soviet Phys. JETP* **24** (1967) 1049–1056.
- [9] F. Calogero, Universal integrable nonlinear PDEs, in: *Application of Analytic and Geometric Methods to Nonlinear Differential Equations* (NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., vol. **413**, P.A. Clarkson, Editor), Kluwer, Dordrecht (1993) 109–114.
- [10] P.A. Clarkson, Painlevé equations – nonlinear special functions, in: *Orthogonal Polynomials and Special Functions: Computation and Application* (Lecture Notes in Math., vol. **1883**, F. Märcellan and W. Van Assche, Editors), Springer-Verlag, Berlin (2006) 331–411.
- [11] P.A. Clarkson, E.L. Mansfield and A.E. Milne, Symmetries and exact solutions of a $(2+1)$ -dimensional sine-Gordon system, *Philos. Trans. Roy. Soc. London Ser. A* **354** (1996) 1807–1835.
- [12] P.A. Clarkson, On Airy solutions of the second Painlevé equation, *Stud. Appl. Math.* **137** (2016) 93–109.
- [13] P.A. Clarkson, Remarks on the Yablonskii-Vorob'ev polynomials, *Phys. Lett. A* **319** (2003) 137–144.
- [14] P.A. Clarkson and E.L. Mansfield, The second Painlevé equation, its hierarchy and associated special polynomials, *Nonlinearity* **16** (2003) R1–R26.
- [15] F.J. Dyson, Dynamics of a spinning gas cloud, *J. Math. Mech.* **18** (1968) 91–101.
- [16] A.S. Fokas and Y.C. Yortsos, The transformation properties of the sixth Painlevé equation and one-parameter families of solutions, *Lett. Nuovo Cimento* **30** (1981) 539–544.
- [17] P.J. Forrester and N.S. Witte, Application of the τ -function theory of Painlevé equations to random matrices: PIV, PII and the GUE, *Commun. Math. Phys.* **219** (2001) 357–398.
- [18] B.D. Fried and Y.H. Ichikawa, On the nonlinear Schrödinger equation for Langmuir waves, *J. Phys. Soc. Japan* **34** (1973) 1073–1082.
- [19] B. Gambier, Sur les équations différentielles du second ordre et du premier degré dont l'intégrale générale est à points critiques fixés, *Acta Math.* **33** (1910) 1–55.
- [20] J.A. Giannini and R.I. Joseph, The role of the second Painlevé transcendent in nonlinear optics, *Phys. Lett. A* **141** (1989) 417–419.

- [21] V.I. Gromak, Bäcklund transformations of Painlevé equations and their applications, in: *The Painlevé Property. One Century Later* (CRM Ser. Math. Phys., R. Conte, Editor), Springer-Verlag, New York (1999) 687–734.
- [22] V.I. Gromak, I. Laine and S. Shimomura, *Painlevé Differential Equations in the Complex Plane* (Studies in Mathematics, vol. **28**), de Gruyter, Berlin (2002).
- [23] Y.H. Ichikawa, T. Imamura and T. Taniuti, Nonlinear wave modulation in collisionless plasma, *J. Phys. Soc. Japan* **33** (1972) 189–197.
- [24] A. Jeffrey, The role of the Korteweg-de Vries equation in plasma physics, *Q. J. L. Roy. Astron. Soc.* **14** (1973) 183–189.
- [25] M. Jimbo and T. Miwa, Monodromy preserving deformations of linear ordinary differential equations with rational coefficients. II, *Physica* **D2** (1981) 407–448.
- [26] T. Kakutani, H. Ono, T. Taniuti and C.C. Wei, Reduction perturbation method in nonlinear wave propagation II: application to hydromagnetic waves in a cold plasma, *J. Phys. Soc. Japan* **24** (1968) 1159–1166.
- [27] T. Kakutani and H. Ono, Weak nonlinear hydromagnetic waves in cold collisionless plasma, *J. Phys. Soc. Japan* **26** (1969) 1305–1318.
- [28] V.I. Karpman, *Nonlinear Waves in Dispersive Media*, Pergamon, Oxford (1975).
- [29] B. Konopelchenko and C. Rogers, On 2+1-dimensional nonlinear systems of Loewner-type, *Phys. Lett. A* **158** (1991) 391–397.
- [30] B. Konopelchenko and C. Rogers, On generalised Loewner systems: novel integrable equations in 2+1-dimensions, *J. Math. Phys.* **34** (1993) 214–242.
- [31] B.G. Konopelchenko, W.K. Schief and C. Rogers, The 2 + 1-dimensional sine Gordon system: its auto Bäcklund transformation, *Phys. Lett. A* **172** (1992) 39–46.
- [32] G.L. Lamb Jr, Analytic descriptions of ultrashort optical pulse propagation in a resonant medium, *Rev. Mod. Phys.* **43** (1971) 99–124.
- [33] J.H. Lee, O.K. Pashaev, C. Rogers and W.K. Schief, The resonant nonlinear Schrödinger equation in cold plasma physics. Application of Bäcklund-Darboux transformations and superposition principles, *J. Plasma Phys.* **73** (2007) 257–272.
- [34] C. Loewner, A transformation theory of partial differential equations of gasdynamics, *NACA Technical Note* **2065** (1950) 1–56.
- [35] C. Loewner, Generation of solutions of systems of partial differential equations by composition of infinitesimal Bäcklund transformations, *J. Anal. Math.* **2** (1952) 219–242.
- [36] S.Y. Lou, C. Rogers and W.K. Schief, Virasoro structure and localised excitations of the LKR system, *J. Math. Phys.* **44** (2003) 5869–5887.
- [37] N.A. Lukashovich, The second Painlevé equation, *Diff. Eq.* **7** (1971) 853–854.
- [38] E. Madelung, Quantentheorie in Hydrodynamischen form, *Zeit. für Phys.* **40** (1926) 322–326.
- [39] A.E. Milne, P.A. Clarkson and A.P. Bassom, Bäcklund transformations and solution hierarchies for the third Painlevé equation, *Stud. Appl. Math.* **98** (1997) 139–194.
- [40] J.J.C. Nimmo, A class of solutions of the Konopelchenko-Rogers equations, *Phys. Lett. A* **168** (1992) 113–119.
- [41] J.J.C. Nimmo and W.K. Schief, Superposition principles associated with the Moutard transformation: an integrable discretisation of a 2 + 1-dimensional sine-Gordon system, *Proc. Roy. Soc. London Ser. A* **453** (1997) 255–279.
- [42] K. Okamoto, Studies on the Painlevé equations III. Second and fourth Painlevé equations, P_{II} and P_{IV} , *Math. Ann.* **275** (1986) 221–255.
- [43] O.K. Pashaev and J.H. Lee, Resonance solitons as black holes in Madelung fluid, *Mod. Phys. Lett. A* **17** (2002) 1601–1619.
- [44] O.K. Pashaev, J.H. Lee and C. Rogers, Soliton resonances in a generalised nonlinear Schrödinger equation, *J. Phys. A: Math. Theor.* **41** (2008) 452001–452009.
- [45] C. Rogers, On the Heisenberg spin equation in hydrodynamics (Informes de Matématics, Série B, vol. **127**) Instituto de Matématica Pura e Applicada, Rio de Janeiro, Brazil (2000).

- [46] C. Rogers and W.K. Schief, On geodesic hydrodynamic motions. Heisenberg spin connections, *J. Math. Anal. Appl.* **251** (2000) 855–870.
- [47] C. Rogers and J.G. Kingston, Non-dissipative magnetohydrodynamic flows with magnetic and velocity field lines orthogonal geodesics on a normal congruence, *SIAM J. Applied Math.* **26** (1974) 183–195.
- [48] C. Rogers and A. Szereszewski, On the geometry of complex-lamellar magnetohydrodynamics. Universal motions, *Stud. Appl. Math.* **26** (2012) 183–195.
- [49] C. Rogers and W.K. Schief, Intrinsic geometry of the NLS equation and its auto-Bäcklund transformation, *Stud. Appl. Math.* **101** (1998) 267–287.
- [50] C. Rogers and W.K. Schief, *Bäcklund and Darboux Transformations. Geometry and Modern Applications in Soliton Theory* (Cambridge Texts in Applied Mathematics), Cambridge University Press, Cambridge (2002).
- [51] C. Rogers and W.K. Schief, Novel integrable reductions in nonlinear continuum mechanics via geometric constraints, *J. Math. Phys.* **44** (2003) 3341–3369.
- [52] C. Rogers and W.K. Schief, Vortex trains in super-Alfvénic magnetogasdynamics. Application of reciprocal-Bäcklund transformations, *J. Nonlinear Math. Phys.* **12** Supplement 1 (2005) 548–564.
- [53] C. Rogers, J.G. Kingston and W.F. Shadwick, On reciprocal-type invariant transformations in magnetogasdynamics, *J. Math. Phys.* **21** (1980) 395–397.
- [54] C. Rogers, A Ermakov-Ray-Reid reduction in 2 + 1-dimensional transverse magnetogasdynamics, in: *Group analysis of differential equations and integrable systems*, Department of Mathematics and Statistics, University of Cyprus, Nicosia (2011) 164–177.
- [55] C. Rogers and W.K. Schief, The pulsodion in 2 + 1-dimensional magnetogasdynamics. Hamiltonian structure and integrability, *J. Math. Phys.* **52** (2011) 083701.
- [56] C. Rogers, B. Malomed, K. Chow and H. An, Ermakov-Ray-Reid systems in nonlinear optics, *J. Phys. A: Math. Theor.* **43** (2010) 455214.
- [57] C. Rogers, B. Malomed and H. An, Ermakov-Ray-Reid reductions of variational approximations in nonlinear optics, *Stud. Appl. Math.* **129** (2012) 389–413.
- [58] C. Rogers and H. An, Ermakov-Ray-Reid systems in 2+1-dimensional rotating shallow water theory, *Stud. Appl. Math.* **125** (2010) 275–299.
- [59] C. Rogers and H. An, A non-isothermal spinning magnetogasdynamics cloud system. Integrable and Hamiltonian-Ermakov structure, *Note di Matematica* **32** (2012) 175–191.
- [60] C. Rogers, Integrable substructure in a Korteweg capillarity model. A Kármán-Tsien type constitutive relation, *J. Nonlinear Math. Phys.* **21** (2014) 74–88.
- [61] C. Rogers, A novel Ermakov-Painlevé II system: n+1-dimensional coupled NLS and elastodynamic reductions, *Stud. Appl. Math.* **133** (2014) 214–231.
- [62] C. Rogers and W.F. Shadwick, *Bäcklund Transformations and Their Applications*, Academic Press, New York (1982).
- [63] C. Rogers, Moving boundary problems for the Harry Dym equation and its reciprocal associates, *Zeit. Angew. Math. Phys.* **66** (2015) 2069–2079.
- [64] C. Rogers, On a class of moving boundary problems for the potential mKdV equation, Special Issue on Waves and Stability, *Ricerche di Matematica* **65** (2016) 563–577.
- [65] C. Rogers, Moving boundary problems for an extended Dym equation. Reciprocal connections, *Mechanica* **52** (2017) 3531–3540.
- [66] W.K. Schief, Hidden integrability in ideal magnetohydrodynamics. The Pohlmeyer-Lund-Regge model, *Phys. Plasmas* **10** (2003) 2677–2685.
- [67] W.K. Schief, On a 2+1-dimensional integrable Ernst-type equation, *Proc. Roy. Soc. London A* **466** (1994) 381–398.
- [68] W.K. Schief, On The geometry of an integrable 2 + 1-dimensional sine-Gordon system, *Proc. Roy. Soc. London Ser. A* **453** (1997) 1671–1688.
- [69] A. Seeger, H. Donth and A. Kochendörfer, Theorie der Versetzungen in eindimensionalen Atomreihen III Versetzungen, Eigenbewegungen und ihr Wechselwirkung, *Zeit. Phys.* **134** (1953) 173–193.
- [70] K. Shimizu and Y.H. Ichikawa, Automodulation of ion oscillation modes in plasma, *J. Phys. Soc. Japan* **33** (1972) 789–792.

- [71] A.P. Vorob'ev, On rational solutions of the second Painlevé equation, *Diff. Eq.* **1** (1965) 79–61.
- [72] H. Washimi and T. Taniuti, Propagation of ion-acoustic solitary waves of small amplitude, *Phys. Rev. Lett.* **17** (1996) 996–998.
- [73] J. Weiland and H. Wilhelmsson, *Coherent Nonlinear Interaction of Waves in Plasmas* (International Series in Natural Philosophy, vol. **88**), Pergamon, New York and Oxford (1997).
- [74] G.B. Whitham, Nonlinear dispersive waves, *Proc. Roy. Soc. London Ser. A* **283** (1965) 238–261.
- [75] A.I. Yablonskii, On rational solutions of the second Painlevé equation, *Vesti. Akad. Nauk. BSSR Ser. Fiz. Tkh. Nauk.* **3** (1959) 30–35.