

A Note on Finite Groups all of Whose Subgroups are C-Normal

Z. Mostaghim

School of Mathematics, Iran University of Science and Technology, Tehran, Iran

Abstract - A subgroup H of a group G is said to be c -normal in G if there exists a normal subgroup N of G such that $HN=G$ and $H \cap N \leq Core(H)$ where $Core(H)$ is the largest normal subgroup of G contained in H . In this paper we consider finite p -groups of order at most p^4 where p is a prime and show that all of their subgroups are c -normal. Also we study some classes of finite groups whose all of subgroups are c -normal.

Index Terms - c -normal subgroups, p -groups, maximal class, supersolvable groups.

I. Introduction

The notion of c -normal subgroup was introduced for the first time by Wang¹. He used the c -normality of maximal subgroups to give some conditions for the solvability and supersolvability of a finite group. For example, he showed that G is solvable if and only if M is c -normal in G for every maximal subgroup M of G . In this paper we consider finite p -groups of order at most p^4 where p is a prime and show that all of their subgroups are c -normal. Also we study some classes of finite groups whose all of subgroups are c -normal.

Throughout, all groups are assumed to be finite groups. Our terminology and notation is standard, see².

II. Preliminaries

In this section, we give some definitions and basic results which are essential in the sequel.

Definition 2.1.¹

Let G be a group. We call a subgroup H c -normal in G if there exists a normal subgroup N of G such that $HN=G$ and $H \cap N \leq Core(H)$.

It is clear that a normal subgroup of G is a c -normal subgroup of G but the converse is not true.

Definition 2.2.¹

We call a group G is c -simple if G has no c -normal subgroup except the identity group 1 and G .

We can easily show that G is c -simple if and only if G is simple.

Lemma 2.3.¹

Let G be a group. Then

- (1) If H is normal in G , then H is c -normal in G ; G is c -simple if and only if G is simple;
- (2) If H is c -normal in G , $H \leq K \leq G$, then H is c -normal in K ;
- (3) Let K be normal in G and $K \leq H$. Then H is c -normal in G if and only if H/K is c -normal in G/K .

Let p be a prime. Now we will give some properties of

non-abelian groups of order p^4 .

Lemma 2.4.³

Let G be a finite non-abelian p -group of order p^4 . Then

- (1) $|Z(G)| = p$ or p^2 ;
- (2) $|G'| \leq p^2$.

Lemma 2.5.³

Let G be a finite non-abelian p -group of order p^4 . If $Z(G)$ is cyclic of order p , then G' has order p^2 . Moreover $Z(G) < G'$ and $G/Z(G)$ is not abelian.

Definition 2.6.⁴

A p -group G is said to be a special p -group if either G is an elementary abelian p -group or we have $\Phi(G) = Z(G) = G'$ and G' is elementary abelian. If the center of a non-abelian special p -group G is cyclic, then G is called extraspecial.

Definition 2.7.⁴

A group of order p^n is said to be a group of maximal class if the class of G is $n - 1$.

Theorem 2.8.⁴

The groups $G = D_{2m}, Q_{2m}, SD_{2m}$ have the following properties.

- (1) The center $Z(G)$ has order 2 and $G/Z(G) \cong D_m$.
- (2) The derived group coincides with $\Phi(G)$ and the class of G is $n - 1$ where $|G| = 2^n$.
- (3) The group Q_{2m} contains exactly one element of order 2.

III. Main Results

Lemma 3.1.

Let G be an extraspecial p -group. Then every subgroup of order p is c -normal.

Proof. It is easy to see that $|G'| = p$. Let H be a subgroup of order p . If $H \cap G' = G'$, then $H \triangleleft G$. By lemma 2.3 H is c -normal in G . If $H \cap G' = 1$, then there exists a maximal subgroup M such that $H \not\leq M$. So $G = HM$ and $H \cap M \leq Core(H)$. Therefore H is c -normal in G .

Theorem 3.2.

Let G be a p -group of order at most p^4 . Then all of subgroups of G are c -normal.

Proof.

If G be an abelian group, then all of subgroups of G are normal and by lemma 2.3, they are c -normal. If G be a non-abelian group of order p^3 , then by lemma 3.1 all of subgroups of order p are c -normal. It is easy to see that other subgroups of G are normal. If G be a non-abelian group of order p^4 , then

by lemmas 2.4 and 2.5 we have two cases:

Case1.

Let $|G'| = p$, therefore $|Z(G)| = p^2$.

Let H be a subgroup of order p . If $H \cap G' \neq 1$, then $H = G'$ and H is a normal subgroup.

Let $H \cap G' = 1$. If there exists a maximal subgroup M such that $H \not\subseteq M$, then $G = HM$ and $H \cap M \leq Core(H)$. Therefore H is a c -normal subgroup.

Let $H \leq \Phi(G)$, so $\Phi(G) = HG'$ and $|\Phi(G)| = p^2$. Since $G/Z(G)$ is an elementary abelian group, so $\Phi(G) \leq Z(G)$ and therefore H is a normal subgroup.

Let H be a subgroup of order p^2 . We have $|H \cap Z(G)| = 1, p$ or p^2 . If $|H \cap Z(G)| = p^2$, then $H = Z(G)$ and H is a normal subgroup. If $|H \cap Z(G)| = 1$, then $G = HZ(G)$ and $H \cap Z(G) \leq Core(H)$. Hence H is c -normal in G .

Let $|H \cap Z(G)| = p$. It is easy to see that $Z(G) = \Phi(G)$. If $H = \Phi(G)$, then H is a normal subgroup. Otherwise there exists a maximal subgroup M such that $HM = G$ and $|H \cap M| = p$. It is easy to see that $H \cap M \leq Core(H)$ and therefore H is a c -normal subgroup.

Case2.

Let $|G'| = p^2$, therefore $|Z(G)| = p$.

Let H be a subgroup of order p . If $H \cap Z(G) = H$, then H is a normal subgroup. Let $H \cap Z(G) = 1$. If there exists a maximal subgroup M such that $H \cap M = 1$, then H is a c -normal subgroup. Otherwise $H \leq \Phi(G) = G' = HZ(G)$. It is easy to see that $H \leq \Phi(G)$ and then H is a normal subgroup.

Let H be a subgroup of order p^2 . Hence $|H \cap G'| = p$ or p^2 . If $|H \cap G'| = p^2$, then H is a normal subgroup.

Let $|H \cap G'| = p$. If $H = \Phi(G)$, then H is a normal subgroup. Otherwise, there exists a maximal subgroup M such that $G = HM$ and $|H \cap M| = p$. Since $H \cap G' \leq Core(H)$, therefore $Core(H) = H \cap G' = H \cap M$. Hence H is a c -normal subgroup.

Theorem 3.3.

Let $G = D_{2n} = \langle a, b \mid a^n = 1, b^2 = 1, bab = a^{-1} \rangle$. Then all of subgroups of G are c -normal.

Proof.

When considered geometrically, D_{2n} consist of n rotations and n reflections of the regular n -gon. The subgroups of D_{2n} are two types:

- (1) Those containing rotations only.
- (2) Those containing rotations and reflections.

Let H be a subgroup of G . We consider two cases.

Case1.

Let H has no reflection. Then $H = \langle a^j \rangle$ for $0 \leq j \leq n - 1$. Thus by lemma 2.3 H is c -normal in G .

Case2.

Let H be of type 2.

- (1) Let $a^j \notin H$ for $0 < j \leq n - 1$, so we have $|H| = 2$. Now let $N = \langle a \rangle$. Then N is a normal subgroup, $G = HN, H \cap N = 1$. Hence H is a c -normal subgroup.

- (2) Let there exists $i > 0$ such that $a^i \in H$. Now let $m = \min\{i \mid i > 0, a^i \in H\}$ and $N = \langle a \rangle$, then $|H| = 2l$, $(1 < l \leq n)$ and $HN = G$. Also we have $H \cap N = \langle a^m \rangle$, then $H \cap N \leq Core(H)$. Hence H is c -normal in G .

Theorem 3.4.

Let G be a 2-group of maximal class. Then all of subgroups of G are c -normal.

Proof.

Since G is a 2-group of maximal class, then G is $D_{2^m} (m \geq 3), Q_{2^m}, SD_{2^m}$. We consider three cases.

Case1.

Let $G = D_{2^m} (m \geq 3)$. Then by theorem 3.3 all of subgroups of G are c -normal.

Case2.

Let $G = Q_m$, then by theorem 2.8 $G/Z(G) \cong D_{2^{m-1}}$. Let H be a subgroup of G such that $Z(G) \subseteq H$, then by lemma 2.3 $H/Z(G)$ is a c -normal subgroup of $G/Z(G)$. By using lemma 2.3 we have H is a c -normal subgroup of G . If H be a subgroup of G and $Z(G) \not\subseteq H$, then $x^j \in H$ for $0 < j \leq 2^{m-1}$ so we have $|H| = 2$. Now let $N = \langle x \rangle$. Thus N is a normal subgroup of G and $G = HN, H \cap N = 1$. Hence H is c -normal in G .

Corollary 3.5.

Let G be one of the following groups.

- (1) A non-nilpotent finite group that all of proper subgroups are Nilpotent.
- (2) A non-abelian finite group that all of proper subgroups are abelian. Then every p -Sylow subgroup of G is c -normal.

Proof.

For case (i) we can see $|G| = p^\alpha q^\beta$, where p and q are distinct primes. Also one of Sylow subgroups of G is cyclic and another is normal. Then every p -Sylow subgroup of G is c -normal. Case (ii) is similar.

Corollary 3.6.

Let G be a finite supersolvable group and $p \mid |G|$ where p is the smallest prime divisor of $|G|$. Then p -Sylow subgroup of G is c -normal.

Proof.

Let $|G| = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_n^{\alpha_n}$, where p_i are primes such that $p_1 > p_2 > \cdots > p_n$. ($p = p_n$) Let P_i be a p_i -Sylow subgroup of G for $1 \leq i \leq n$, then $P_1 P_2 \cdots P_k$ is a normal subgroup for all $1 \leq k \leq n$. It is easy to see that P_n is c -normal in G .

IV. GAP Program

In this section we use GAP⁵ and give a program for finding c -normal subgroups. By using this program we can find all c -normal subgroups of a finite group with two generations. With a few changes in this program we can find a program for finding c -normal subgroups in a finite group with any number of generations and relations.

F:=FreeGroup("a","b");

```

a:=GeneratorsOfGroup(F)[1];
b:=GeneratorsOfGroup(F)[2];
Read("r");
G:=F/r;
n:=Order(G);
z:=LowIndexSubgroupsFpGroup(G,TrivialSubgroup(G),n);
s:=[];
  for i in [1..Size(z)] do
    t:=ConjugacyClassSubgroups(G,z[i]);
    for j in [1..Size(t)] do
      Add(s,t[j]);
    od;
  od;
cnorm:=[];
N:=[];
H:=[];
  for i in [1..Size(s)] do
    vi:=IsNormal(G,s[i]);
    if vi=true then Add(N,s[i]);fi;
    if vi=false then Add(H,s[i]);fi;
  od;
for y in [1..Size(H)] do
l:=0;
m:=0;

```

```

h:=false;
while (m=0 or h=false) and l<=Size(N) do
l:=l+1;
eH:=Elements(H[y]);
eN:=Elements(N[l]);
HN:=[];
  for i in [1..Order(H[y])] do
    for j in [1..Order(N[l])] do
      u:=eH[i]*eN[j];
      AddSet(HN,u);
    od;
  od;
h:=IsSubgroup(Core(G,H[y]),Intersection(H[y],N[l]));
if HN=G then m:=1;fi;
  od;
if HN=G and h=true then Add(cnorm,H[y]);fi;
od;

```

References And Notes

- [1] Y. Wang , J.Algebra 180, (1996).
- [2] D. Gorenstein, Finite groups,Chelsea Publishing Company (1980).
- [3] H.Behravesh and H.Mousavi ,Proc.Indian Acad.Sci(Math. Sci) 119, (2009).
- [4] M. Suzuki, Group Theory II, Springer-Verlag,New York, (1986).
- [5] The GAPGroup,GAP-Groups,Algorithms and Programming, (2005).