

A Class of Nonlinear Stochastic Volatility Models

Jun Yu, Zhenlin Yang

Singapore Management University

Abstract

This paper proposes a class of nonlinear stochastic volatility (SV) models based on the Box-Cox transformation. The proposed class encompasses many parametric SV models that have appeared in the literature, including the well known lognormal SV model, and has an advantage in the ease with which different specifications on SV can be tested. In addition, the functional form of transformation which induces marginal normality of volatility is obtained as a byproduct of this general way of modeling SV. Efficient method of moments is used to estimate the model. Empirical results reveal that the lognormal SV model is rejected.

Keywords: Box-Cox Transform; EMM; GARCH

1. Introduction

Modeling the volatility of financial time series via stochastic volatility (SV) models has received a great deal of attention both in the theoretical finance and in the empirical finance literature. Prices of options based on SV models are shown to be more accurate than those based on the Black-Scholes model (see, for example, Melino and Turnbull (1990)). Moreover, the SV model is proved to be a serious competitor to GARCH-type models for explaining time varying volatility. Empirical successes of the lognormal SV model relative to GARCH are documented in Kim, et al. (1998) in terms of in-sample fitting, and in Yu (2002) in terms of out-of-sample forecasting.

In the option pricing literature, the SV model is often formulated in terms of stochastic differential equations. For instance, Wiggins (1987), Chesney and Scott (1989), and Scott (1987) specify the following model for the asset price $P(t)$ and the corresponding volatility $\sigma^2(t)$,

$$dP(t)/P(t) = \alpha dt + \sigma(t)dB_1(t), \quad (0.1)$$

$$d \ln \sigma^2(t) = \lambda(\xi - \ln \sigma^2(t))dt + \gamma dB_2(t), \quad (0.2)$$

where $B_1(t)$ and $B_2(t)$ are two independent standard Brownian motions.

In the empirical literature, the above continuous time model is often discretized. For example, the Euler-Maruyama The approximation, after a location shift and

reparameterization, leads to the so-called *lognormal* SV model given by

$$X_t = \sigma_t e_t, \quad (0.3)$$

$$\ln \sigma_t^2 = \mu + \phi(\ln \sigma_{t-1}^2 - \mu) + \sigma v_t, \quad (0.4)$$

where X_t is the continuously compounded return and e_t, v_t are two uncorrelated sequences of independent and identically distributed (iid) $N(0, 1)$ variables. In the literature σ_t is often represented by $\exp(\frac{1}{2}h_t)$. One implication of its specification is that the marginal distribution of logarithmic volatility is normal. This assumption has very important implications for financial economics and risk management. So far this model as received a great deal of attention in the econometrics literature.

Alternative SV models have appeared in the literature. For example, Stein and Stein (1991) and Johnson and Shanno (1987) assume $\sigma(t)$ follows, respectively, an Ornstein-Uhlenbeck (OU) process and a geometric Brownian motion, while Hull and White (1987) and Heston (1993) assume a geometric Brownian motion and a square-root process for $\sigma^2(t)$. In the discrete time case, various SV models can be regarded as generalizations to corresponding GARCH models. For example, a polynomial SV model is a generalization of GARCH(1,1) (Bollerslev (1986)) while a square root polynomial SV model is a generalization of standard deviation GARCH(1,1). Andersen (1994) introduces a general class of SV models, of which a class of polynomial SV models has been emphasized. This class encompasses many discrete time SV models in the literature.

Despite all these alternative specifications, there is a lack of procedure for selecting appropriate functional form of SV. The specification of the correct SV function, on the other hand, is very important in two respects. First, different functional forms lead to different formulae for option pricing. Misspecification of the SV function can result in incorrect option prices. Second, the marginal distribution of volatility depends upon the functional form of SV.

In this paper, we propose a new class of SV models, namely, the nonlinear SV model. Like the class of Andersen (1994), it includes as special cases many discrete time SV models that have appeared in the literature. It overlaps with but does not encompass the class of Andersen.

An advantage of our proposed class is the ease with which different specifications on SV can be tested. In fact, the specification test is based on a single parameter. Another advantage of our proposed class is that, as a byproduct of this general way of modeling volatility, one obtains the functional form of transformation which induces marginal normality of volatility. Section 2 presents this class of nonlinear models. In Section 3, we use efficient method of moments (EMM) to estimate the model. In Section 4, the class is fitted to daily observations on an index return.

2. A Class of Nonlinear SV Models

The lognormal SV model specifies that the logarithmic volatility follows an AR(1) process. However, this relationship may not always be warranted by the data. A natural generalization to this relationship is to allow a general (nonlinear) smooth function of volatility to follow an AR(1) process. That is,

$$X_t = \sigma_t e_t, \quad (0.5)$$

$$h(\sigma_t^2, \delta) = \mu + \phi[h(\sigma_{t-1}^2, \delta) - \mu] + \sigma v_t, \quad (0.6)$$

where e_t and v_t are two uncorrelated $N(0, 1)$ sequences, and h is a smooth function indexed by a parameter δ . A nice choice of this function is the Box-Cox power function (Box and Cox (1964)):

$$h(t, \delta) = \begin{cases} (t^\delta - 1)/\delta, & \text{if } \delta \neq 0, \\ \log t, & \text{if } \delta = 0. \end{cases} \quad (0.7)$$

As the function h is specified as a general nonlinear function, we call the model the *nonlinear* SV model. Several attractive features of this new class of SV models include: i) as we will show below it includes the lognormal SV model and the other ‘‘classical’’ SV models as special cases, ii) it adds great flexibility on the functional form, and iii) it allows a simple test for the lognormal SV specification, i.e., a test of $H_0 : \delta = 0$, and some other ‘‘classical’’ SV specifications. If we write $h_t = h(\sigma_t^2, \delta)$, then we can re-write the nonlinear SV models as

$$X_t = [g(h_t, \delta)]^{1/2} e_t, \quad (0.8)$$

$$h_t = \mu + \phi(h_{t-1} - \mu) + \sigma v_t, \quad (0.9)$$

where $g(h_t, \delta)$ is the inverse Box-Cox transformation:

$$g(h_t, \delta) = \begin{cases} (1 + \delta h_t)^{1/\delta}, & \text{if } \delta \neq 0, \\ \exp(h_t), & \text{if } \delta = 0. \end{cases} \quad (0.10)$$

Equivalently we can re-write them in a form of

$$X_t = \sigma_t e_t, \quad (0.11)$$

$$\frac{(\sigma_t^2)^\delta - 1}{\delta} = \mu + \phi\left[\frac{(\sigma_{t-1}^2)^\delta - 1}{\delta} - \mu\right] + \sigma v_t. \quad (0.12)$$

Table 1: Alternative Stochastic Volatility Models and Parameter Relationship

| | Models | δ | μ | ϕ |
|--------------------------------|---|----------|-------|--------|
| Wiggins Scott Kim et al. | $\ln \sigma_t^2 = \mu +$ $\phi(\ln \sigma_{t-1}^2 - \mu) + \sigma v_t$ | 0 | | |
| Scott Andersen | $\sigma_t = \mu +$ $\phi(\sigma_{t-1} - \mu) + \sigma v_t$ | 0.5 | | |
| Heston | $\sigma_t = \phi \sigma_{t-1} + \sigma v_t$ | 0.5 | 0 | |
| Hull & White | $\ln \sigma_t^2 = \mu +$ $\ln \sigma_{t-1}^2 + \sigma v_t$ | 0 | | 1 |
| Andersen | $\sigma_t^2 = \mu +$ $\phi(\sigma_{t-1}^2 - \mu) + \sigma v_t$ | 1 | | |
| Clark | $\ln \sigma_t^2 = \mu + \sigma v_t$ | 0 | | 0 |
| Nonlinear SV | $\frac{((\sigma_t^2)^\delta - 1)/\delta = \mu +$ $\phi[(\sigma_{t-1}^2)^\delta - 1]/\delta - \mu]$ $+ \sigma v_t$ | | | |

Denote the parameters of interest by $\theta = (\mu, \delta, \phi, \sigma)$.

The idea of our proposed SV models is similar to that in Higgins and Bera (1992) from the linear ARCH model (Engle (1982)) to the nonlinear ARCH (NARCH) model. Obviously, our model provides a stochastic volatility generalization of a nonlinear GARCH(1,1) model.

It can be seen as $\delta \rightarrow 0$, $(1 + \delta h_t)^{1/(2\delta)} \rightarrow \exp(0.5 h_t)$ and $((\sigma_t^2)^\delta - 1)/\delta \rightarrow \ln \sigma_t^2$. Hence the proposed nonlinear SV model includes the lognormal SV model as a special case. If $\delta = 1$, the variance equation (0.12) becomes

$$\sigma_t^2 = \mu' + \phi(\sigma_{t-1}^2 - \mu') + \sigma v_t, \quad (0.13)$$

where $\mu' = \mu + 1$. This is a polynomial SV model in Andersen (1994). According to this specification, volatility follows a normal distribution as its marginal distribution. If $\delta = 0.5$, the variance equation (0.12) becomes

$$\sigma_t = \mu'' + \phi(\sigma_{t-1} - \mu'') + 0.5 \sigma v_t, \quad (0.14)$$

where $\mu'' = 0.5\mu + 1$. This is a square root polynomial SV model in Andersen (1994) and can be regarded as a discrete time version of the continuous time SV model in Scott (1987) and Stein and Stein (1991). As a result, the marginal distribution of the square root of volatility is Gaussian.

In Table 1 we summarize some well-known stochastic volatility models and show their parameter relations with our model. For the continuous time stochastic volatility models, their Euler discrete time versions are considered. It can be seen that all these models can be obtained from

our model by placing the appropriate restrictions on the three parameters δ, μ and ϕ . In fact, all the models except ours require δ to be 0, 0.5, or 1.¹ For a general δ , our model is different from any of them and δ provides some idea about the degree of departure from a “classical” parametric SV model.

The Box-Cox transformation has been applied in various areas in finance. One of the most relevant applications to our work may be that by Higgins and Bera (1992) who introduce the nonlinear ARCH model. Another relevant application is Hentschel (1995) who introduces a family of GARCH models by applying the Box-Cox transformation to the conditional standard deviation. A nice feature of our proposed class is that it provides a simple way to test the null hypothesis of polynomial SV specifications, including the lognormal SV specification, against a variety of non-polynomial alternatives. In fact, this specification test is based entirely on a single parameter, δ . Moreover, as a consequence of specification testing, our proposed class provides an effective channel to check the marginal distribution of unobserved volatility. Therefore, our method serves as an alternative approach for studying marginal distribution of daily volatility to the approach based on ultra-high frequency data that has appeared in a recent literature (cf Andersen et al., 2001).

3. Estimation by EMM

In this paper we use EMM to estimate the proposed model and to make inferences. EMM is first introduced by Gallant and Tauchen (1996) and has now found many successful applications in economics and finance; see Gallant and Tauchen (2001a) for a brief review of the literature. It is closely related to GMM of Hansen (1982). An important difference between them is that while GMM relies on an *ad hoc* chosen set of moment conditions, EMM is based on a judiciously chosen set of moment conditions. The moment conditions EMM employs is the expectation of the score of an auxiliary model which is often referred to as the score generator.

Let the SV model be the structural model. The conditional density of the structural model is defined by

$$p_t(x_t|y_t, \theta),$$

where y_t is a vector of lagged x_t , the true value of θ is θ_0 and $\theta_0 \in \Theta \subset \mathcal{R}^{\ell_\theta}$ with ℓ_θ being the length of θ_0 . Denote the conditional density of an auxiliary model by

$$f_t(x_t|y_t, \beta), \beta \in R \subset \mathcal{R}^{\ell_\beta}.$$

Further define the expected score of the auxiliary model under the structural model as

$$m(\theta, \beta) = \int \cdots \int \frac{\partial}{\partial \beta} \ln f(x|y, \beta) p(x|y, \theta) p(y|\theta) dx dy.$$

Obviously, in the context of the SV model, the integration cannot be solved analytically since neither $p(x|y, \theta)$ nor $p(y|\theta)$ has closed form. However, it is easy to simulate from an SV model so that one can approximate the integral by Monte Carlo simulations. That is

$$m(\theta, \beta) \approx m_N(\theta, \beta) \equiv \frac{1}{N} \sum_{\tau=1}^N \frac{\partial}{\partial \beta} \ln f(\hat{x}_\tau(\theta) | \hat{y}_\tau(\theta), \beta),$$

where $\{\hat{x}_\tau, \hat{y}_\tau\}$ are simulated from the structural model. The EMM estimator is a minimum chi-squared estimator which minimizes the following quadratic form,

$$\hat{\theta}_n = \arg \min_{\theta \in \Theta} m'_N(\theta, \hat{\beta}_n) (I_n)^{-1} m_N(\theta, \hat{\beta}_n),$$

where $\hat{\beta}_n$ is a quasi maximum likelihood estimator of the auxiliary model and I_n is an estimate of

$$I_0 = \lim_{n \rightarrow \infty} \text{Var} \left(\frac{1}{\sqrt{n}} \sum_{t=1}^n \left\{ \frac{\partial}{\partial \beta} \ln f_t(x_t | y_t, \beta^*) \right\} \right)$$

with β^* being the pseudo true value of β . Under regularity conditions, Gallant and Tauchen (1996) show that the EMM estimator is consistent and has the following asymptotic normal distribution,

$$\sqrt{n}(\hat{\theta}_n - \theta_0) \xrightarrow{d} N(0, \frac{\partial}{\partial \theta} m(\theta_0, \beta^*) (I_0)^{-1} \frac{\partial}{\partial \theta'} m(\theta_0, \beta^*)).$$

For specification testing, we have

$$J_n = n m'_N(\hat{\theta}_n, \hat{\beta}_n) (I_n)^{-1} m_N(\hat{\theta}_n, \hat{\beta}_n) \xrightarrow{d} \chi_{\ell_\beta - \ell_\theta}^2$$

under the null hypothesis that the structural model is correct. When a model fails the above specification test one may wish to examine the quasi-t-ratios and/or t-ratios to look for some suggestion as to what is wrong with the structural model. Large quasi-t-ratios and t-ratios reveal the features of the data that the structural model cannot approximate.

Furthermore, Gallant and Tauchen (1996) show that if the auxiliary model nests the data generating process, EMM has the same asymptotic variance as maximum likelihood and hence is fully efficient. If the auxiliary model closely approximate the data generating process, EMM is nearly fully efficient (Gallant and Long (1997)).

To choose an auxiliary model, the seminonparametric (SNP) density proposed by Gallant and Tauchen (1989) can be used since its success has been documented in many applications. As to SNP modeling, six out of eight tuning

¹Some specifications in Table 1 may be different from the actual specifications used in the original references. However, they are equivalent to each other via Ito's lemma. For example, Heston (1993) adopts a square root specification for σ_t^2 which is identical to assuming σ_t follows a particular OU process.

parameters are to be selected, namely, $L_u, L_g, L_r, L_p, K_z,$ and K_y . The other two parameters, I_z and I_x , are irrelevant for univariate time series and hence are set to 0. L_u determines the location transformation whereas L_g and L_r determine the scale transformation. Altogether they determine the nature of the leading term of the Hermite expansion. The other two parameters K_z and K_y determine the nature of the innovation. To search for a good auxiliary model, one can use the Schwarz BIC criterion to move along an upward expansion path until an adequate model is found, as outlined in Bansal et al. (1995). To preserve space we refer readers to Gallant and Tauchen (2001b) for further discussion about the role of the tuning parameters and how to design an expansion path to choose them.

4. An Empirical Application

In this section we consider an application using a stock index series which consist of 4044 observations on 100 times the log-first difference of the daily S&P 500 index for the period from January 4, 1977 to December 31, 1992. The same data have been used in Gallant and Tauchen (2001a) to fit the SV model of Clark (1973). Since the return series is not mean-adjusted, to allow for a possible no-zero mean and also some dynamics in mean, we introduce an AR(1) structure in the mean equation. As a consequence, we fit the following two models to the data:

$$X_t = \mu_0 + c(X_{t-1} - \mu_0) + \exp\left(\frac{1}{2}h_t\right)e_t, \quad (0.15)$$

$$h_t = \mu + \phi(h_{t-1} - \mu) + \sigma v_t; \quad (0.16)$$

and

$$X_t = \mu_0 + c(X_{t-1} - \mu_0) + (1 + \delta h_t)^{1/(2\delta)} e_t, \quad (0.17)$$

$$h_t = \mu + \phi(h_{t-1} - \mu) + \sigma v_t. \quad (0.18)$$

We call them the lognormal SV model and the proposed SV model respectively.

The same sets of tuning parameters as in Gallant and Tauchen (2001a) are employed in the SNP model, since the identical dataset is used. We report these tuning parameters in the following order

$$(L_u, L_g, L_r, L_p, K_z, I_z, K_y, I_y).$$

To ensure that the chosen SNP model is reasonable, we have compared the BIC value with those from many alternative sets of tuning parameters and find that the BIC value from the chosen SNP model is one of the smallest. The set of the tuning parameters, the corresponding BIC value, the leading term in the Hermite expansion, the characterization of X_t and the number of parameters in the auxiliary model are presented in Table 2 where an ARCH leading term is used.

Table 2: Tuning parameters for SNP modeling, BIC, Leading Term, Characterization of X_t and the Number of Parameters in the Auxiliary Model

| Data | Tuning parameters | BIC |
|--------------|---------------------------|--------------|
| SP500 | (2, 0, 11, 1, 4, 0, 0, 0) | 1.32715 |
| Leading term | Characterization of X_t | ℓ_β |
| ARCH | Semiparametric ARCH | 19 |

Since the sample size is large, we believe that the choice of leading term is not crucial as long as a form of conditional heteroskedasticity has been accommodated. In a Monte Carlo study, Andersen et al. (1999) find that the EMM efficiency approaches that of maximum likelihood for larger sample size when various forms of conditional heteroskedasticity are used as the leading term. Moreover, they find that the EMM-based inferences, such as the t -statistic and J_n statistic, are robust to the choice of auxiliary model when the sample size is large.

Table 3 and Table 4 report the empirical results, where we perturb starting values when doing the optimizations. Furthermore, we simulate 101,000 observations from the SV models, of which first 1,000 observations are discarded in order to let transients die out. Table 3 reports the estimates, the numerical Wald standard errors, the 95% approximate criterion-difference confidence intervals, the value of statistic J_n , and the degrees of freedom and the p -value of J_n for the lognormal SV model and for the proposed SV model. Table 4 reports the quasi- t -ratios and t -ratios from the score generator for both models.

A few results emerge from these two tables. First and most interestingly, the point estimate of δ is -0.4597 . It is significantly less than 0 and hence significantly less than 0.5 and 1. As a consequence, one has to reject the lognormal SV model and all the other SV models in Table 1. Observing that δ is not significantly different from -0.5 , to gain some idea about our estimated results, we approximate $\delta \approx -0.5$, plug the estimates into Equation (0.12) and get the following estimated variance equation:

$$\frac{1}{\sigma_t^2} = 1.2237 + 0.9840\left(\frac{1}{\sigma_{t-1}^2} - 1.2237\right) + 0.057v_t.$$

This compares to the estimated variance equation in the lognormal model,

$$\ln \sigma_t^2 = -0.3425 + 0.9846(\ln \sigma_{t-1}^2 + 0.3425) + 0.1022v_t.$$

It can be easily demonstrated that the two marginal distributions are not close to each other and hence the lognormal distribution is not a good approximation to the

Table 3: Parameter estimates, standard errors, confidence intervals, χ^2 criterion for S&P500

| | Lognormal SV model | Proposed SV model |
|-------------|---|---|
| μ_0 | 0.0389 (0.0136) [0.0123, 0.0651] | 0.0387 (0.0137) [0.0122, 0.0650] |
| c | 0.0880 (0.0159) [0.0583, 0.1179] | 0.0875 (0.0158) [0.0579, 0.1177] |
| μ | -0.3425 (0.0613) [-0.4593, -0.2291] | -0.4474 (0.0812) [-0.5066, -0.3441] |
| ϕ | 0.9846 (0.0120) [0.9657, 0.9966] | 0.9840 (0.0106) [0.9736, 0.9946] |
| σ | 0.1022 (0.0456) [0.0566, 0.1719] | 0.1140 (0.0405) [0.0734, 0.1532] |
| δ | NA | -0.4597 (0.1807) [-0.6316, -0.2885] |
| χ^2 | 25.01 | 21.65 |
| df | 14 | 13 |
| p - value | 0.034 | 0.061 |

marginal distribution of volatility. Furthermore, theoretically speaking a distributional constraint has to be imposed on h_t in the proposed SV models to ensure non-negativeness of σ_t . In the empirical applications, however, we still adopt the assumption of exact normality. To understand how restricted this assumption is, we calculate $\text{Prob}(\sigma_t < 0) = \text{Prob}(1/\sigma_t < 0) = 0.0000065$ which is a very small value. Besides, our model is not limited to the Box-Cox power transform. Some other transformations, which do not have this restriction (such as the one in Yang (2006)), may be used.

Second, the point estimate of ϕ (0.984) is close to 1 and just in the stationary region. It remains almost the same in the proposed model. In fact all the estimated parameters have similar magnitude and standard errors across both models. The only exceptions are μ which decreases from -0.3425 to -0.4474 and σ which increases from 0.1022 to 0.1140. This is because μ and σ are closely related to δ in the proposed model. Since the estimated δ is far away from 0 in the proposed model, this translates

Table 4: Quasi-t-ratios and t-ratios for S&P500

| | | Lognormal SV model | | Proposed SV model | |
|------|----------|--------------------|---------|-------------------|---------|
| | | Quasi-t-ratio | T-ratio | Quasi-t-ratio | T-ratio |
| VAR | b_1 | -0.096 | -0.334 | -0.089 | -0.302 |
| | b_2 | -1.243 | -1.249 | -1.243 | -1.259 |
| | b_3 | -0.446 | -1.540 | -0.436 | -1.571 |
| ARCH | r_1 | 0.781 | 1.391 | 0.676 | 1.471 |
| | r_2 | 0.627 | 0.909 | 0.513 | 0.896 |
| | r_3 | 0.130 | 0.224 | 0.028 | 0.051 |
| | r_4 | 1.122 | 1.228 | 1.047 | 1.160 |
| | r_5 | 1.053 | 1.498 | 0.933 | 1.399 |
| | r_6 | 0.253 | 0.312 | 0.132 | 0.179 |
| | r_7 | 0.066 | 0.088 | -0.052 | -0.071 |
| | r_8 | 1.032 | 1.521 | 0.861 | 1.489 |
| | r_9 | 0.881 | 1.040 | 0.754 | 0.945 |
| | r_{10} | 0.549 | 0.588 | 0.448 | 0.494 |
| | r_{11} | -0.243 | -0.280 | -0.388 | -0.500 |
| | r_{12} | 0.257 | 0.270 | 0.145 | 0.159 |
| SNP | s_1 | 0.033 | 0.063 | 0.093 | 0.179 |
| | s_2 | 0.977 | 3.091 | 0.706 | 2.477 |
| | s_3 | 0.205 | 0.232 | 0.258 | 0.291 |
| | s_4 | 2.303 | 3.767 | 1.798 | 3.332 |

to large discrepancies between the estimated μ 's and the estimated σ 's. Third, the minimum χ^2 criterion provides mild evidence against the lognormal specification, which is rejected at the 5% level but not at the 1% level. The evidence is consistent with the diagnostic quasi-t-ratios and t-ratios. There are large quasi-t-ratios and t-ratios on the scores corresponding to the polynomial part of the SNP score. These t-statistics indicate that $\exp(0.5h_t)$ may not be the correct transformation. When the proposed SV model is fitted, the p -value of J_n statistic increases by about 80%. One can accept the proposed model at the 5% level. Furthermore, all the quasi-t-ratios become insignificant in the proposed model. Although some of the t-ratios on the scores corresponding to the polynomial part of the SNP score are still too large, they are clearly smaller than those in the lognormal model.

References

- [1] Andersen, T. (1994). Stochastic autoregressive volatility: A framework for volatility modeling. *Math. Finance* 4, 75–102.
- [2] Andersen, T., T. Bollerslev, F.X. Diebold, and H. Ebens (2001). The distribution of realized stock re-

- turn volatility. *J. of Financial Economics* 61, 43–76.
- [3] Andersen, T., H. Chung and B. Sorensen (1999). Efficient method of moments estimation of a stochastic volatility model: A Monte Carlo study. *J. of Econometrics* 91, 61–87
- [4] Bansal, R., A.R. Gallant, R. Hussey and G. Tauchen (1995) Nonparametric estimation of structural models for high-frequency currency market data. *J. of Econometrics* 66, 251–287.
- [5] Bollerslev, T. (1986) Generalized autoregressive conditional heteroskedasticity. *J. of Econometrics* 31, 307–327.
- [6] Box, G.E.P. and D.R. Cox (1964). An analysis of transformations. *J. Roy. Stat. Soc. Ser. B* 26, 211–243.
- [7] Chesney, M. and L.O. Scott (1989). Pricing European currency options: A comparison of the modified Black-Scholes model and a random variance model. *J. of Financial and Quantitative Analysis* 24, 267–284.
- [8] Clark, P.K. (1973). A subordinated stochastic process model with finite variance for speculative price. *Econometrica* 41, 135–155.
- [9] Engle, R.F. (1982) Autoregressive conditional heteroscedasticity with estimates of the variance of United Kingdom inflation. *Econometrica* 50, 987–1007.
- [10] Gallant, A.R. and L.R. Long (1997). Estimating stochastic differential equations efficiently by minimum chi-squared. *Biometrika* 84, 125–141.
- [11] Gallant, A.R. and G. Tauchen (1989). Semiparametric estimation of conditionally constrained heterogeneous processes: Asset pricing applications. *Econometrica* 57, 1091–1120.
- [12] Gallant, A.R. and G. Tauchen (1996). Which moments to match? *Econometric Theory* 12, 657–681.
- [13] Gallant, A.R. and G. Tauchen (2001a). EMM: A program for efficient method of moments estimation. User’s Guide, Department of Economics, University of North Carolina.
- [14] Gallant, A.R. and G. Tauchen (2001b). SNP: A program for nonparametric time series analysis. User’s Guide, Department of Economics, University of North Carolina.
- [15] Hansen, L.P. (1982). Large sample properties of generalized method of moments estimators. *Econometrica* 50, 1029–1054.
- [16] Hentschel, L. (1995). All in the family: Nesting symmetric and asymmetric GARCH models. *J. Financial Economics* 39, 71–104.
- [17] Heston, S.L. (1993). A closed-form solution for options with stochastic volatility, with application to bond and currency options. *Rev. of Financial Stud.* 6, 327–343.
- [18] Higgins, M.L. and A. Bera (1992). A class of nonlinear ARCH models. *International Economic Rev.* 33, 137–158.
- [19] Hull, J. and A. White (1987). The pricing of options on assets with stochastic volatilities. *J. of Finance* 42, 281–300.
- [20] Johnson, H. and D. Shanno (1987). Option pricing when the variance is changing. *J. of Financial and Quantitative Analysis* 22, 143–152.
- [21] Kim, S., N. Shephard, and S. Chib (1998). Stochastic volatility: Likelihood inference and comparison with ARCH models. *Rev. of Economic Stud.* 65, 361–393.
- [22] Melino, A. and S.M. Turnbull (1990). Pricing foreign currency options with stochastic volatility. *J. of Econometrics* 45, 239–265.
- [23] Scott, L.O. (1987). Option pricing when the variance changes randomly: Theory, estimation and an application. *J. of Financial and Quantitative Analysis* 22, 419–439.
- [24] Stein, E.M. and J.C. Stein (1991). Stock price distributions with stochastic volatility: An analytical approach. *Rev. of Financial Stud.* 4, 727–752.
- [25] Wiggins, J.B. (1987) Option values under stochastic volatility: Theory and empirical estimate. *J. of Financial Economics* 19, 351–372.
- [26] Yang, Z. (2006) A Modified Family of Power Transformations. *Economics Letters* 92, 14–19.
- [27] Yu, J. (2002) Forecasting volatility in the New Zealand stock market. *Applied Financial Economics* 12, 193–202.